

Existence of finitely presented intersection-saturated groups

Enric Ventura

Departament de Matemàtiques
Universitat Politècnica de Catalunya

Geometry Seminar

Instytut Matematyczny, Uniwersytet Wrocławski

(joint work with J. Delgado and M. Roy)

June 5th, 2023.

Outline

- 1 Our main results
- 2 Free-times-free-abelian groups
- 3 Realizable / unrealizable k -configurations
- 4 The free case
- 5 Open questions

Outline

- 1 Our main results
- 2 Free-times-free-abelian groups
- 3 Realizable / unrealizable k -configurations
- 4 The free case
- 5 Open questions

Free groups

It is well known that subgroups of free groups are free ...

$$H \leq \mathbb{F}_n \Rightarrow H \text{ is free}$$


but not necessarily of rank $\leq n$.

Example

Consider $\mathbb{F}_2 = \langle x, y \mid \rangle$ and the normal closure of x ,

$$\langle\langle x \rangle\rangle = \langle \dots, y^2xy^{-2}, yxy^{-1}, x, y^{-1}xy, y^{-2}xy^2, \dots \rangle.$$

Looking at its Stallings graph



we see these generators are a free basis; so, $\mathbb{F}_{\aleph_0} \leq \mathbb{F}_2$.

Free groups

It is well known that subgroups of free groups are free ...

$$H \leq \mathbb{F}_n \Rightarrow H \text{ is free}$$


but not necessarily of rank $\leq n$.

Example

Consider $\mathbb{F}_2 = \langle x, y \mid \rangle$ and the normal closure of x ,

$$\langle\langle x \rangle\rangle = \langle \dots, y^2xy^{-2}, yxy^{-1}, x, y^{-1}xy, y^{-2}xy^2, \dots \rangle.$$

Looking at its Stallings graph



we see these generators are a free basis; so, $\mathbb{F}_{\aleph_0} \leq \mathbb{F}_2$.

Free groups

It is well known that subgroups of free groups are free ...

$$H \leq \mathbb{F}_n \Rightarrow H \text{ is free}$$


but not necessarily of rank $\leq n$.

Example

Consider $\mathbb{F}_2 = \langle x, y \mid \rangle$ and the normal closure of x ,

$$\langle\langle x \rangle\rangle = \langle \dots, y^2xy^{-2}, yxy^{-1}, x, y^{-1}xy, y^{-2}xy^2, \dots \rangle.$$

Looking at its Stallings graph



we see these generators are a free basis; so, $\mathbb{F}_{\aleph_0} \leq \mathbb{F}_2$.

Free groups

It is well known that subgroups of free groups are free ...

$$H \leq \mathbb{F}_n \Rightarrow H \text{ is free}$$


but not necessarily of rank $\leq n$.

Example

Consider $\mathbb{F}_2 = \langle x, y \mid \rangle$ and the normal closure of x ,

$$\langle\langle x \rangle\rangle = \langle \dots, y^2xy^{-2}, yxy^{-1}, x, y^{-1}xy, y^{-2}xy^2, \dots \rangle.$$

Looking at its Stallings graph



we see these generators are a free basis; so, $\mathbb{F}_{\aleph_0} \leq \mathbb{F}_2$.

The Howson property

Definition

A group G is **Howson** if, for any finitely generated $H, K \leq_{fg} G$, the intersection $H \cap K$ is, again, finitely generated.

Theorem (Howson, 1954)

Free groups are Howson.

In other words... the configuration



is not realizable in a free group (\circ means f.g. and \bullet means non-f.g.).

Observation

Out of $2^3 = 8$ possible such configurations this is the only one forbidden in free groups.

The Howson property

Definition

A group G is **Howson** if, for any finitely generated $H, K \leq_{fg} G$, the intersection $H \cap K$ is, again, finitely generated.

Theorem (Howson, 1954)

Free groups are Howson.

In other words... the configuration



is not realizable in a free group (\circ means f.g. and \bullet means non-f.g.).

Observation

Out of $2^3 = 8$ possible such configurations this is the only one forbidden in free groups.

The Howson property

Definition

A group G is **Howson** if, for any finitely generated $H, K \leq_{fg} G$, the intersection $H \cap K$ is, again, finitely generated.

Theorem (Howson, 1954)

Free groups are Howson.

In other words... the configuration



is not realizable in a free group (○ means f.g. and ● means non-f.g.).

Observation

Out of $2^3 = 8$ possible such configurations this is the only one forbidden in free groups.

The Howson property

Definition

A group G is *Howson* if, for any finitely generated $H, K \leq_{fg} G$, the intersection $H \cap K$ is, again, finitely generated.

Theorem (Howson, 1954)

Free groups are Howson.

In other words... the configuration



is not realizable in a free group (○ means f.g. and ● means non-f.g.).

Observation




Out of $2^3 = 8$ possible such configurations this is the only one forbidden in free groups.

Intersection configurations

Question

What about configurations with $k \geq 2$ subgroups (k -configurations)?

Using this convention, what about the following 3-configurations?






Intersection configurations

Question

What about configurations with $k \geq 2$ subgroups (k -configurations)?

Using this convention, what about the following 3-configurations?



Our main results

Theorem (Delgado–Roy–V., '22)

A k -configuration is realizable in \mathbb{F}_n , $n \geq 2$, \Leftrightarrow it respects the Howson property.

Theorem (Delgado–Roy–V. '22)

There exist finitely presented intersection-saturated groups.

Our main results

Theorem (Delgado–Roy–V., '22)

A k -configuration is realizable in \mathbb{F}_n , $n \geq 2$, \Leftrightarrow it respects the Howson property.

Theorem (Delgado–Roy–V. '22)

There exist finitely presented intersection-saturated groups.

Formal definitions

Definition

A (*intersection*) k -configuration is a map $\chi: \mathcal{P}([k]) \setminus \{\emptyset\} \rightarrow \{0, 1\}$. If $\mathcal{I} = (1)\chi^{-1}$ is the support of χ , we write $\chi = \chi_{\mathcal{I}}$. Notation:

- $\mathbf{0} = \chi_{\emptyset}$ is the *zero-configuration*;
- $\mathbf{1} = \chi_{\mathcal{P}([k]) \setminus \{\emptyset\}}$ is the *one-configuration*;
- $\chi_{\mathcal{I}}$ is an *almost-zero* k -configuration if $\mathcal{I} = \{I\}$.

Definition

A k -configuration χ is *realizable in a group* G if there exists subgroups $H_1, \dots, H_k \leq G$ such that, for every $\emptyset \neq I \subseteq [k]$, $H_I = \bigcap_{i \in I} H_i$ if f.g. $\Leftrightarrow (I)\chi = 0$. Note that $H_{I \cup J} = H_I \cap H_J$.

Definition

A group G is *intersection-saturated* if every k -configuration is realizable in G .

Formal definitions

Definition

A (*intersection*) k -configuration is a map $\chi: \mathcal{P}([k]) \setminus \{\emptyset\} \rightarrow \{0, 1\}$. If $\mathcal{I} = (1)\chi^{-1}$ is the support of χ , we write $\chi = \chi_{\mathcal{I}}$. Notation:

- $\mathbf{0} = \chi_{\emptyset}$ is the *zero-configuration*;
- $\mathbf{1} = \chi_{\mathcal{P}([k]) \setminus \{\emptyset\}}$ is the *one-configuration*;
- $\chi_{\mathcal{I}}$ is an *almost-zero k -configuration* if $\mathcal{I} = \{I\}$.

Definition

A k -configuration χ is *realizable in a group G* if there exists subgroups $H_1, \dots, H_k \leq G$ such that, for every $\emptyset \neq I \subseteq [k]$, $H_I = \bigcap_{i \in I} H_i$ if f.g. $\Leftrightarrow (I)\chi = 0$. Note that $H_{I \cup J} = H_I \cap H_J$.

Definition

A group G is *intersection-saturated* if every k -configuration is realizable in G .

Formal definitions

Definition

A (*intersection*) k -configuration is a map $\chi: \mathcal{P}([k]) \setminus \{\emptyset\} \rightarrow \{0, 1\}$. If $\mathcal{I} = (1)\chi^{-1}$ is the support of χ , we write $\chi = \chi_{\mathcal{I}}$. Notation:

- $\mathbf{0} = \chi_{\emptyset}$ is the *zero-configuration*;
- $\mathbf{1} = \chi_{\mathcal{P}([k]) \setminus \{\emptyset\}}$ is the *one-configuration*;
- $\chi_{\mathcal{I}}$ is an *almost-zero* k -configuration if $\mathcal{I} = \{I\}$.

Definition

A k -configuration χ is *realizable in a group* G if there exists subgroups $H_1, \dots, H_k \leq G$ such that, for every $\emptyset \neq I \subseteq [k]$, $H_I = \bigcap_{i \in I} H_i$ if f.g. $\Leftrightarrow (I)\chi = 0$. Note that $H_{I \cup J} = H_I \cap H_J$.

Definition

A group G is *intersection-saturated* if every k -configuration is realizable in G .

Formal definitions

Definition

A (*intersection*) k -configuration is a map $\chi: \mathcal{P}([k]) \setminus \{\emptyset\} \rightarrow \{0, 1\}$. If $\mathcal{I} = (1)\chi^{-1}$ is the support of χ , we write $\chi = \chi_{\mathcal{I}}$. Notation:

- $\mathbf{0} = \chi_{\emptyset}$ is the *zero-configuration*;
- $\mathbf{1} = \chi_{\mathcal{P}([k]) \setminus \{\emptyset\}}$ is the *one-configuration*;
- $\chi_{\mathcal{I}}$ is an *almost-zero* k -configuration if $\mathcal{I} = \{I\}$.

Definition

A k -configuration χ is *realizable in a group* G if there exists subgroups $H_1, \dots, H_k \leq G$ such that, for every $\emptyset \neq I \subseteq [k]$, $H_I = \bigcap_{i \in I} H_i$ if f.g. $\Leftrightarrow (I)\chi = 0$. Note that $H_{I \cup J} = H_I \cap H_J$.

Definition

A group G is *intersection-saturated* if every k -configuration is realizable in G .

Formal definitions

Definition

A (*intersection*) k -configuration is a map $\chi: \mathcal{P}([k]) \setminus \{\emptyset\} \rightarrow \{0, 1\}$. If $\mathcal{I} = (1)\chi^{-1}$ is the support of χ , we write $\chi = \chi_{\mathcal{I}}$. Notation:

- $\mathbf{0} = \chi_{\emptyset}$ is the *zero-configuration*;
- $\mathbf{1} = \chi_{\mathcal{P}([k]) \setminus \{\emptyset\}}$ is the *one-configuration*;
- $\chi_{\mathcal{I}}$ is an *almost-zero* k -configuration if $\mathcal{I} = \{I\}$.

Definition

A k -configuration χ is *realizable in a group* G if there exists subgroups $H_1, \dots, H_k \leq G$ such that, for every $\emptyset \neq I \subseteq [k]$, $H_I = \bigcap_{i \in I} H_i$ if f.g. $\Leftrightarrow (I)\chi = \mathbf{0}$. Note that $H_{I \cup J} = H_I \cap H_J$.

Definition

A group G is *intersection-saturated* if every k -configuration is realizable in G .

Formal definitions

Definition

A (*intersection*) k -configuration is a map $\chi: \mathcal{P}([k]) \setminus \{\emptyset\} \rightarrow \{0, 1\}$. If $\mathcal{I} = (1)\chi^{-1}$ is the support of χ , we write $\chi = \chi_{\mathcal{I}}$. Notation:

- $\mathbf{0} = \chi_{\emptyset}$ is the *zero-configuration*;
- $\mathbf{1} = \chi_{\mathcal{P}([k]) \setminus \{\emptyset\}}$ is the *one-configuration*;
- $\chi_{\mathcal{I}}$ is an *almost-zero* k -configuration if $\mathcal{I} = \{I\}$.

Definition

A k -configuration χ is *realizable in a group* G if there exists subgroups $H_1, \dots, H_k \leq G$ such that, for every $\emptyset \neq I \subseteq [k]$, $H_I = \bigcap_{i \in I} H_i$ if f.g. $\Leftrightarrow (I)\chi = \mathbf{0}$. Note that $H_{I \cup J} = H_I \cap H_J$.

Definition

A group G is *intersection-saturated* if every k -configuration is realizable in G .

Outline

- 1 Our main results
- 2 Free-times-free-abelian groups**
- 3 Realizable / unrealizable k -configurations
- 4 The free case
- 5 Open questions

Free-times-free-abelian groups

$$\mathbb{G} = \mathbb{F}_n \times \mathbb{Z}^m = \langle x_1, \dots, x_n, t_1, \dots, t_m \mid [x_i, t_j] = 1, [t_i, t_k] = 1 \rangle.$$

Normal form: $\forall g \in \mathbb{G}, g = w(x_1, \dots, x_n)t_1^{a_1} \cdots t_m^{a_m} = wt^{\mathbf{a}}$, where $\mathbf{a} = (a_1, \dots, a_m) \in \mathbb{Z}^m$. This way, $(ut^{\mathbf{a}})(vt) = uvt^{\mathbf{a}+\mathbf{b}}$.

Observation

These groups sit in a split short exact sequence; and, for $H \leq \mathbb{G}$,

$$\begin{aligned} 1 \rightarrow \mathbb{Z}^m \hookrightarrow \mathbb{G} \xrightarrow{\pi} \mathbb{F}_n \rightarrow 1, \\ 1 \rightarrow L_H = H \cap \mathbb{Z}^m \hookrightarrow H \rightarrow H\pi \rightarrow 1. \end{aligned}$$

Moreover, H is finitely generated $\Leftrightarrow H\pi$ is so.

Free-times-free-abelian groups

$$\mathbb{G} = \mathbb{F}_n \times \mathbb{Z}^m = \langle x_1, \dots, x_n, t_1, \dots, t_m \mid [x_i, t_j] = 1, [t_i, t_k] = 1 \rangle.$$

Normal form: $\forall g \in \mathbb{G}, g = w(x_1, \dots, x_n)t_1^{a_1} \dots t_m^{a_m} = wt^{\mathbf{a}}$, where $\mathbf{a} = (a_1, \dots, a_m) \in \mathbb{Z}^m$. This way, $(ut^{\mathbf{a}})(vt) = uvt^{\mathbf{a}+\mathbf{b}}$.

Observation

These groups sit in a split short exact sequence; and, for $H \leq \mathbb{G}$,

$$\begin{aligned} 1 \rightarrow \mathbb{Z}^m \hookrightarrow \mathbb{G} \xrightarrow{\pi} \mathbb{F}_n \rightarrow 1, \\ 1 \rightarrow L_H = H \cap \mathbb{Z}^m \hookrightarrow H \rightarrow H\pi \rightarrow 1. \end{aligned}$$

Moreover, H is finitely generated $\Leftrightarrow H\pi$ is so.

Free-times-free-abelian groups

$$\mathbb{G} = \mathbb{F}_n \times \mathbb{Z}^m = \langle x_1, \dots, x_n, t_1, \dots, t_m \mid [x_i, t_j] = 1, [t_i, t_k] = 1 \rangle.$$

Normal form: $\forall g \in \mathbb{G}, g = w(x_1, \dots, x_n)t_1^{a_1} \dots t_m^{a_m} = wt^{\mathbf{a}}$, where $\mathbf{a} = (a_1, \dots, a_m) \in \mathbb{Z}^m$. This way, $(ut^{\mathbf{a}})(vt) = uvt^{\mathbf{a}+\mathbf{b}}$.

Observation

These groups sit in a split short exact sequence; and, for $H \leq \mathbb{G}$,

$$\begin{aligned} 1 \rightarrow \mathbb{Z}^m \xrightarrow{\iota} \mathbb{G} \xrightarrow{\pi} \mathbb{F}_n \rightarrow 1, \\ 1 \rightarrow L_H = H \cap \mathbb{Z}^m \hookrightarrow H \rightarrow H\pi \rightarrow 1. \end{aligned}$$

Moreover, H is finitely generated $\Leftrightarrow H\pi$ is so.

Free-times-free-abelian groups

$$\mathbb{G} = \mathbb{F}_n \times \mathbb{Z}^m = \langle x_1, \dots, x_n, t_1, \dots, t_m \mid [x_i, t_j] = 1, [t_i, t_k] = 1 \rangle.$$

Normal form: $\forall g \in \mathbb{G}, g = w(x_1, \dots, x_n)t_1^{a_1} \dots t_m^{a_m} = wt^{\mathbf{a}}$, where $\mathbf{a} = (a_1, \dots, a_m) \in \mathbb{Z}^m$. This way, $(ut^{\mathbf{a}})(vt) = uvt^{\mathbf{a}+\mathbf{b}}$.

Observation

These groups sit in a split short exact sequence; and, for $H \leq \mathbb{G}$,

$$\begin{aligned} 1 \rightarrow \mathbb{Z}^m \xrightarrow{\iota} \mathbb{G} \xrightarrow{\pi} \mathbb{F}_n \rightarrow 1, \\ 1 \rightarrow L_H = H \cap \mathbb{Z}^m \hookrightarrow H \rightarrow H\pi \rightarrow 1. \end{aligned}$$

Moreover, H is finitely generated $\Leftrightarrow H\pi$ is so.

Free-times-free-abelian groups

Proposition (Delgado–V. '13)

Every subgroup $H \leq \mathbb{G}$ admits a (computable) basis

$$H = \langle u_1 t^{\mathbf{a}_1}, u_2 t^{\mathbf{a}_2}, \dots, u_r t^{\mathbf{a}_r}; t^{\mathbf{b}_1}, \dots, t^{\mathbf{b}_s} \rangle,$$

where $\{u_1, \dots, u_r\}$ is a free-basis for $H\pi$, $\mathbf{a}_1, \dots, \mathbf{a}_r \in \mathbb{Z}^m$, $0 \leq r \leq \infty$, $\mathbf{b}_1, \dots, \mathbf{b}_s \in \mathbb{Z}^m$ is an abelian-basis for $L_H = H \cap \mathbb{Z}^m$, and $0 \leq s \leq m$.

Proposition (Moldavanski)

The groups $F_n \times \mathbb{Z}^m$, $n \geq 2$, $m \geq 1$, are not Howson.

Question

Are them intersection-saturated?... no... but collectively yes ...

Theorem (Delgado–Roy–V. '22)

- The set of configs realizable in $\mathbb{F}_n \times \mathbb{Z}^m$ increases strictly with m ;
- Every configuration is realizable in $\mathbb{F}_n \times \mathbb{Z}^m$ for $m \gg 0$.

Free-times-free-abelian groups

Proposition (Delgado–V. '13)

Every subgroup $H \leq \mathbb{G}$ admits a (computable) basis

$$H = \langle u_1 t^{\mathbf{a}_1}, u_2 t^{\mathbf{a}_2}, \dots, u_r t^{\mathbf{a}_r}; t^{\mathbf{b}_1}, \dots, t^{\mathbf{b}_s} \rangle,$$

where $\{u_1, \dots, u_r\}$ is a free-basis for $H\pi$, $\mathbf{a}_1, \dots, \mathbf{a}_r \in \mathbb{Z}^m$, $0 \leq r \leq \infty$, $\mathbf{b}_1, \dots, \mathbf{b}_s \in \mathbb{Z}^m$ is an abelian-basis for $L_H = H \cap \mathbb{Z}^m$, and $0 \leq s \leq m$.

Proposition (Moldavanski)

The groups $F_n \times \mathbb{Z}^m$, $n \geq 2$, $m \geq 1$, **are not** Howson.

Question

Are them intersection-saturated?... no... but collectively yes ...

Theorem (Delgado–Roy–V. '22)

- The set of configs realizable in $\mathbb{F}_n \times \mathbb{Z}^m$ increases strictly with m ;
- Every configuration is realizable in $\mathbb{F}_n \times \mathbb{Z}^m$ for $m \gg 0$.

Free-times-free-abelian groups

Proposition (Delgado–V. '13)

Every subgroup $H \leq \mathbb{G}$ admits a (computable) basis

$$H = \langle u_1 t^{\mathbf{a}_1}, u_2 t^{\mathbf{a}_2}, \dots, u_r t^{\mathbf{a}_r}; t^{\mathbf{b}_1}, \dots, t^{\mathbf{b}_s} \rangle,$$

where $\{u_1, \dots, u_r\}$ is a free-basis for $H\pi$, $\mathbf{a}_1, \dots, \mathbf{a}_r \in \mathbb{Z}^m$, $0 \leq r \leq \infty$, $\mathbf{b}_1, \dots, \mathbf{b}_s \in \mathbb{Z}^m$ is an abelian-basis for $L_H = H \cap \mathbb{Z}^m$, and $0 \leq s \leq m$.

Proposition (Moldavanski)

The groups $F_n \times \mathbb{Z}^m$, $n \geq 2$, $m \geq 1$, **are not** Howson.

Question

Are them intersection-saturated?... no... but collectively yes ...

Theorem (Delgado–Roy–V. '22)

- The set of configs realizable in $\mathbb{F}_n \times \mathbb{Z}^m$ increases strictly with m ;
- Every configuration is realizable in $\mathbb{F}_n \times \mathbb{Z}^m$ for $m \gg 0$.

Free-times-free-abelian groups

Proposition (Delgado–V. '13)

Every subgroup $H \leq \mathbb{G}$ admits a (computable) basis

$$H = \langle u_1 t^{\mathbf{a}_1}, u_2 t^{\mathbf{a}_2}, \dots, u_r t^{\mathbf{a}_r}; t^{\mathbf{b}_1}, \dots, t^{\mathbf{b}_s} \rangle,$$

where $\{u_1, \dots, u_r\}$ is a free-basis for $H\pi$, $\mathbf{a}_1, \dots, \mathbf{a}_r \in \mathbb{Z}^m$, $0 \leq r \leq \infty$, $\mathbf{b}_1, \dots, \mathbf{b}_s \in \mathbb{Z}^m$ is an abelian-basis for $L_H = H \cap \mathbb{Z}^m$, and $0 \leq s \leq m$.

Proposition (Moldavanski)

The groups $F_n \times \mathbb{Z}^m$, $n \geq 2$, $m \geq 1$, **are not** Howson.

Question

Are them intersection-saturated?... ... no... but collectively yes ...

Theorem (Delgado–Roy–V. '22)

- The set of configs realizable in $\mathbb{F}_n \times \mathbb{Z}^m$ increases strictly with m ;
- Every configuration is realizable in $\mathbb{F}_n \times \mathbb{Z}^m$ for $m \gg 0$.

Free-times-free-abelian groups

Proposition (Delgado–V. '13)

Every subgroup $H \leq \mathbb{G}$ admits a (computable) basis

$$H = \langle u_1 t^{\mathbf{a}_1}, u_2 t^{\mathbf{a}_2}, \dots, u_r t^{\mathbf{a}_r}; t^{\mathbf{b}_1}, \dots, t^{\mathbf{b}_s} \rangle,$$

where $\{u_1, \dots, u_r\}$ is a free-basis for $H\pi$, $\mathbf{a}_1, \dots, \mathbf{a}_r \in \mathbb{Z}^m$, $0 \leq r \leq \infty$, $\mathbf{b}_1, \dots, \mathbf{b}_s \in \mathbb{Z}^m$ is an abelian-basis for $L_H = H \cap \mathbb{Z}^m$, and $0 \leq s \leq m$.

Proposition (Moldavanski)

The groups $F_n \times \mathbb{Z}^m$, $n \geq 2$, $m \geq 1$, **are not** Howson.

Question

Are them intersection-saturated?... ... no... but collectively yes ...

Theorem (Delgado–Roy–V. '22)

- The set of configs realizable in $\mathbb{F}_n \times \mathbb{Z}^m$ increases strictly with m ;
- Every configuration is realizable in $\mathbb{F}_n \times \mathbb{Z}^m$ for $m \gg 0$.

Free-times-free-abelian groups

Proposition (Delgado–V. '13)

Every subgroup $H \leq \mathbb{G}$ admits a (computable) basis

$$H = \langle u_1 t^{\mathbf{a}_1}, u_2 t^{\mathbf{a}_2}, \dots, u_r t^{\mathbf{a}_r}; t^{\mathbf{b}_1}, \dots, t^{\mathbf{b}_s} \rangle,$$

where $\{u_1, \dots, u_r\}$ is a free-basis for $H\pi$, $\mathbf{a}_1, \dots, \mathbf{a}_r \in \mathbb{Z}^m$, $0 \leq r \leq \infty$, $\mathbf{b}_1, \dots, \mathbf{b}_s \in \mathbb{Z}^m$ is an abelian-basis for $L_H = H \cap \mathbb{Z}^m$, and $0 \leq s \leq m$.

Proposition (Moldavanski)

The groups $F_n \times \mathbb{Z}^m$, $n \geq 2$, $m \geq 1$, **are not** Howson.

Question

Are them intersection-saturated?... ... no... but collectively yes ...

Theorem (Delgado–Roy–V. '22)

- The set of configs realizable in $\mathbb{F}_n \times \mathbb{Z}^m$ increases strictly with m ;
- Every configuration is realizable in $\mathbb{F}_n \times \mathbb{Z}^m$ for $m \gg 0$.

Free-times-free-abelian groups

Proposition (Delgado–V. '13)

Every subgroup $H \leq \mathbb{G}$ admits a (computable) basis

$$H = \langle u_1 t^{\mathbf{a}_1}, u_2 t^{\mathbf{a}_2}, \dots, u_r t^{\mathbf{a}_r}; t^{\mathbf{b}_1}, \dots, t^{\mathbf{b}_s} \rangle,$$

where $\{u_1, \dots, u_r\}$ is a free-basis for $H\pi$, $\mathbf{a}_1, \dots, \mathbf{a}_r \in \mathbb{Z}^m$, $0 \leq r \leq \infty$, $\mathbf{b}_1, \dots, \mathbf{b}_s \in \mathbb{Z}^m$ is an abelian-basis for $L_H = H \cap \mathbb{Z}^m$, and $0 \leq s \leq m$.

Proposition (Moldavanski)

The groups $F_n \times \mathbb{Z}^m$, $n \geq 2$, $m \geq 1$, **are not** Howson.

Question

Are them intersection-saturated?... ... no... but collectively yes ...

Theorem (Delgado–Roy–V. '22)

- The set of configs realizable in $\mathbb{F}_n \times \mathbb{Z}^m$ increases strictly with m ;
- Every configuration is realizable in $\mathbb{F}_n \times \mathbb{Z}^m$ for $m \gg 0$.

Free-times-free-abelian groups

Theorem (Delgado–V. '13)

There is an algorithm which, on input (a set of generators for) $H, K \leq_{fg} \mathbb{G}$, decides whether $H \cap K$ is f.g. and, if so, computes a basis for it.

(Sketch of proof)

Given (basis for) subgroups $H_1, H_2 \leq_{fg} \mathbb{G} = \mathbb{F}_n \times \mathbb{Z}^m$, consider

$$\begin{array}{ccccc}
 & & (H_1 \cap H_2)\pi & & \\
 & & \triangle & & \\
 H_1\pi & \xleftarrow{i_1} & H_1\pi \cap H_2\pi & \xleftarrow{i_2} & H_2\pi \\
 \downarrow \rho_1 & \quad \quad \quad & \downarrow \rho & \quad \quad \quad & \downarrow \rho_2 \\
 \mathbb{Z}^{r_1} & \xleftarrow{P_1} & \mathbb{Z}^r & \xrightarrow{P_2} & \mathbb{Z}^{r_2} \\
 & \swarrow A_1 & \downarrow R & \searrow A_2 & \\
 & & \mathbb{Z}^m & & \\
 & & \vee & & \\
 & & L_1 + L_2 & &
 \end{array}$$

A calculation shows that $(H_1 \cap H_2)\pi = (L_1 + L_2)R^{-1}\rho^{-1} \trianglelefteq H_1\pi \cap H_2\pi$.

So, $H_1 \cap H_2$ is f.g. $\Leftrightarrow \begin{cases} r = 0, 1 \text{ or} \\ r \geq 2 \text{ and } (H_1 \cap H_2)\pi \leq_{fi} H_1\pi \cap H_2\pi. \end{cases}$ □


Free-times-free-abelian groups

Theorem (Delgado–V. '13)

There is an algorithm which, on input (a set of generators for) $H, K \leq_{fg} \mathbb{G}$, decides whether $H \cap K$ is f.g. and, if so, computes a basis for it.

(Sketch of proof)

Given (basis for) subgroups $H_1, H_2 \leq_{fg} \mathbb{G} = \mathbb{F}_n \times \mathbb{Z}^m$, consider



A calculation shows that $(H_1 \cap H_2)\pi = (L_1 + L_2)R^{-1}\rho^{-1} \trianglelefteq H_1\pi \cap H_2\pi$.

So, $H_1 \cap H_2$ is f.g. $\Leftrightarrow \begin{cases} r = 0, 1 \text{ or} \\ r \geq 2 \text{ and } (H_1 \cap H_2)\pi \leq_{fi} H_1\pi \cap H_2\pi. \end{cases}$ □

Free-times-free-abelian groups

Theorem (Delgado–Roy–V. '22)

There is an algorithm which, on input (a set of generators for) $H_1, \dots, H_k \leq_{fg} \mathbb{G}$, decides whether $H_1 \cap \dots \cap H_k$ is f.g. and, if so, computes a basis for it.

Proposition

*Let $M', M'' \leq \mathbb{F}_n$ be such that $\langle M', M'' \rangle = M' * M''$. Then, for any $H'_1, \dots, H'_k \leq M' \leq \mathbb{F}_n$ and $H''_1, \dots, H''_k \leq M'' \leq \mathbb{F}_n$,*

$$\bigcap_{j=1}^k \langle H'_j, H''_j \rangle = \left\langle \bigcap_{j=1}^k H'_j, \bigcap_{j=1}^k H''_j \right\rangle.$$

Free-times-free-abelian groups

Theorem (Delgado–Roy–V. '22)

There is an algorithm which, on input (a set of generators for) $H_1, \dots, H_k \leq_{fg} \mathbb{G}$, decides whether $H_1 \cap \dots \cap H_k$ is f.g. and, if so, computes a basis for it.

Proposition

*Let $M', M'' \leq \mathbb{F}_n$ be such that $\langle M', M'' \rangle = M' * M''$. Then, for any $H'_1, \dots, H'_k \leq M' \leq \mathbb{F}_n$ and $H''_1, \dots, H''_k \leq M'' \leq \mathbb{F}_n$,*

$$\bigcap_{j=1}^k \langle H'_j, H''_j \rangle = \left\langle \bigcap_{j=1}^k H'_j, \bigcap_{j=1}^k H''_j \right\rangle.$$

Free-times-free-abelian groups

Observation

The same is **not true** in $\mathbb{G} = \mathbb{F}_n \times \mathbb{Z}^m$, even with $M', M'' \leq \mathbb{G}$ in strongly complementary position, i.e., $\langle M'\pi, M''\pi \rangle = M'\pi * M''\pi$ and $\langle M'\tau, M''\tau \rangle = M'\tau \oplus M''\tau$.

Example

Consider $\mathbb{G} = \mathbb{F}_4 \times \mathbb{Z}^2 = \langle x_1, x_2, x_3, x_4 \mid - \rangle \times \langle t_1, t_2 \mid [t_1, t_2] \rangle$,
 $M' = \langle x_1, x_2, t^{(1,0)} \rangle$, $M'' = \langle x_3, x_4, t^{(0,1)} \rangle$, and the respective subgroups

- $H'_1 = \langle x_1, x_2 \rangle$, $H'_2 = \langle x_1 t^{(1,0)}, x_2 \rangle \leq M'$, and
- $H''_1 = \langle x_3, x_4 \rangle$, $H''_2 = \langle x_3 t^{(0,1)}, x_4 \rangle \leq M''$.

We have $H'_1 \cap H'_2 = \langle x_1^{-i} x_2 x_1^i, i \in \mathbb{Z} \rangle$, $H''_1 \cap H''_2 = \langle x_3^{-i} x_4 x_3^i, i \in \mathbb{Z} \rangle$, and

$\langle H'_1 \cap H'_2, H''_1 \cap H''_2 \rangle = (H'_1 \cap H'_2) * (H''_1 \cap H''_2) = \langle x_1^{-i} x_2 x_1^i, x_3^{-i} x_4 x_3^i \mid i \in \mathbb{Z} \rangle$,

which **does not contain** $x_3^{-1} x_2 x_3 \in \langle H'_1, H''_1 \rangle = \langle x_1, x_2, x_3, x_4 \rangle$
 $\in \langle H'_2, H''_2 \rangle = \langle x_1 t^{(1,0)}, x_2, x_3 t^{(0,1)}, x_4 \rangle$.

Free-times-free-abelian groups

Observation

The same is **not true** in $\mathbb{G} = \mathbb{F}_n \times \mathbb{Z}^m$, even with $M', M'' \leq \mathbb{G}$ in strongly complementary position, i.e., $\langle M'\pi, M''\pi \rangle = M'\pi * M''\pi$ and $\langle M'\tau, M''\tau \rangle = M'\tau \oplus M''\tau$.

Example

Consider $\mathbb{G} = \mathbb{F}_4 \times \mathbb{Z}^2 = \langle x_1, x_2, x_3, x_4 \mid - \rangle \times \langle t_1, t_2 \mid [t_1, t_2] \rangle$,
 $M' = \langle x_1, x_2, t^{(1,0)} \rangle$, $M'' = \langle x_3, x_4, t^{(0,1)} \rangle$, and the respective subgroups

- $H'_1 = \langle x_1, x_2 \rangle$, $H'_2 = \langle x_1 t^{(1,0)}, x_2 \rangle \leq M'$, and
- $H''_1 = \langle x_3, x_4 \rangle$, $H''_2 = \langle x_3 t^{(0,1)}, x_4 \rangle \leq M''$.

We have $H'_1 \cap H'_2 = \langle x_1^{-i} x_2 x_1^i, i \in \mathbb{Z} \rangle$, $H''_1 \cap H''_2 = \langle x_3^{-i} x_4 x_3^i, i \in \mathbb{Z} \rangle$, and

$\langle H'_1 \cap H'_2, H''_1 \cap H''_2 \rangle = (H'_1 \cap H'_2) * (H''_1 \cap H''_2) = \langle x_1^{-i} x_2 x_1^i, x_3^{-i} x_4 x_3^i \mid i \in \mathbb{Z} \rangle$,

which **does not contain** $x_3^{-1} x_2 x_3 \in \langle H'_1, H''_1 \rangle = \langle x_1, x_2, x_3, x_4 \rangle$
 $\in \langle H'_2, H''_2 \rangle = \langle x_1 t^{(1,0)}, x_2, x_3 t^{(0,1)}, x_4 \rangle$.

Free-times-free-abelian groups

Observation

The same is **not true** in $\mathbb{G} = \mathbb{F}_n \times \mathbb{Z}^m$, even with $M', M'' \leq \mathbb{G}$ in strongly complementary position, i.e., $\langle M'\pi, M''\pi \rangle = M'\pi * M''\pi$ and $\langle M'\tau, M''\tau \rangle = M'\tau \oplus M''\tau$.

Example

Consider $\mathbb{G} = \mathbb{F}_4 \times \mathbb{Z}^2 = \langle x_1, x_2, x_3, x_4 \mid - \rangle \times \langle t_1, t_2 \mid [t_1, t_2] \rangle$,
 $M' = \langle x_1, x_2, t^{(1,0)} \rangle$, $M'' = \langle x_3, x_4, t^{(0,1)} \rangle$, and the respective subgroups

- $H'_1 = \langle x_1, x_2 \rangle$, $H'_2 = \langle x_1 t^{(1,0)}, x_2 \rangle \leq M'$, and
- $H''_1 = \langle x_3, x_4 \rangle$, $H''_2 = \langle x_3 t^{(0,1)}, x_4 \rangle \leq M''$.

We have $H'_1 \cap H'_2 = \langle x_1^{-i} x_2 x_1^i, i \in \mathbb{Z} \rangle$, $H''_1 \cap H''_2 = \langle x_3^{-i} x_4 x_3^i, i \in \mathbb{Z} \rangle$, and

$\langle H'_1 \cap H'_2, H''_1 \cap H''_2 \rangle = (H'_1 \cap H'_2) * (H''_1 \cap H''_2) = \langle x_1^{-i} x_2 x_1^i, x_3^{-i} x_4 x_3^i \mid i \in \mathbb{Z} \rangle$,

which **does not contain** $x_3^{-1} x_2 x_3 \in \langle H'_1, H''_1 \rangle = \langle x_1, x_2, x_3, x_4 \rangle$
 $\in \langle H'_2, H''_2 \rangle = \langle x_1 t^{(1,0)}, x_2, x_3 t^{(0,1)}, x_4 \rangle$.

Free-times-free-abelian groups

Observation

The same is **not true** in $\mathbb{G} = \mathbb{F}_n \times \mathbb{Z}^m$, even with $M', M'' \leq \mathbb{G}$ in strongly complementary position, i.e., $\langle M'\pi, M''\pi \rangle = M'\pi * M''\pi$ and $\langle M'\tau, M''\tau \rangle = M'\tau \oplus M''\tau$.

Example

Consider $\mathbb{G} = \mathbb{F}_4 \times \mathbb{Z}^2 = \langle x_1, x_2, x_3, x_4 \mid - \rangle \times \langle t_1, t_2 \mid [t_1, t_2] \rangle$,
 $M' = \langle x_1, x_2, t^{(1,0)} \rangle$, $M'' = \langle x_3, x_4, t^{(0,1)} \rangle$, and the respective subgroups

- $H'_1 = \langle x_1, x_2 \rangle$, $H'_2 = \langle x_1 t^{(1,0)}, x_2 \rangle \leq M'$, and
- $H''_1 = \langle x_3, x_4 \rangle$, $H''_2 = \langle x_3 t^{(0,1)}, x_4 \rangle \leq M''$.

We have $H'_1 \cap H'_2 = \langle x_1^{-i} x_2 x_1^i, i \in \mathbb{Z} \rangle$, $H''_1 \cap H''_2 = \langle x_3^{-i} x_4 x_3^i, i \in \mathbb{Z} \rangle$, and

$\langle H'_1 \cap H'_2, H''_1 \cap H''_2 \rangle = (H'_1 \cap H'_2) * (H''_1 \cap H''_2) = \langle x_1^{-i} x_2 x_1^i, x_3^{-i} x_4 x_3^i \mid i \in \mathbb{Z} \rangle$,

which **does not contain** $x_3^{-1} x_2 x_3 \in \langle H'_1, H''_1 \rangle = \langle x_1, x_2, x_3, x_4 \rangle$
 $\in \langle H'_2, H''_2 \rangle = \langle x_1 t^{(1,0)}, x_2, x_3 t^{(0,1)}, x_4 \rangle$.

Free-times-free-abelian groups

Observation

The same is **not true** in $\mathbb{G} = \mathbb{F}_n \times \mathbb{Z}^m$, even with $M', M'' \leq \mathbb{G}$ in strongly complementary position, i.e., $\langle M'\pi, M''\pi \rangle = M'\pi * M''\pi$ and $\langle M'\tau, M''\tau \rangle = M'\tau \oplus M''\tau$.

Example

Consider $\mathbb{G} = \mathbb{F}_4 \times \mathbb{Z}^2 = \langle x_1, x_2, x_3, x_4 \mid - \rangle \times \langle t_1, t_2 \mid [t_1, t_2] \rangle$,
 $M' = \langle x_1, x_2, t^{(1,0)} \rangle$, $M'' = \langle x_3, x_4, t^{(0,1)} \rangle$, and the respective subgroups

- $H'_1 = \langle x_1, x_2 \rangle$, $H'_2 = \langle x_1 t^{(1,0)}, x_2 \rangle \leq M'$, and
- $H''_1 = \langle x_3, x_4 \rangle$, $H''_2 = \langle x_3 t^{(0,1)}, x_4 \rangle \leq M''$.

We have $H'_1 \cap H'_2 = \langle x_1^{-i} x_2 x_1^i, i \in \mathbb{Z} \rangle$, $H''_1 \cap H''_2 = \langle x_3^{-i} x_4 x_3^i, i \in \mathbb{Z} \rangle$, and

$\langle H'_1 \cap H'_2, H''_1 \cap H''_2 \rangle = (H'_1 \cap H'_2) * (H''_1 \cap H''_2) = \langle x_1^{-i} x_2 x_1^i, x_3^{-i} x_4 x_3^i \mid i \in \mathbb{Z} \rangle$,

which **does not contain** $x_3^{-1} x_2 x_3 \in \langle H'_1, H''_1 \rangle = \langle x_1, x_2, x_3, x_4 \rangle$
 $\in \langle H'_2, H''_2 \rangle = \langle x_1 t^{(1,0)}, x_2, x_3 t^{(0,1)}, x_4 \rangle$.

Free-times-free-abelian groups

Observation

The same is **not true** in $\mathbb{G} = \mathbb{F}_n \times \mathbb{Z}^m$, even with $M', M'' \leq \mathbb{G}$ in strongly complementary position, i.e., $\langle M'\pi, M''\pi \rangle = M'\pi * M''\pi$ and $\langle M'\tau, M''\tau \rangle = M'\tau \oplus M''\tau$.

Example

Consider $\mathbb{G} = \mathbb{F}_4 \times \mathbb{Z}^2 = \langle x_1, x_2, x_3, x_4 \mid - \rangle \times \langle t_1, t_2 \mid [t_1, t_2] \rangle$,
 $M' = \langle x_1, x_2, t^{(1,0)} \rangle$, $M'' = \langle x_3, x_4, t^{(0,1)} \rangle$, and the respective subgroups

- $H'_1 = \langle x_1, x_2 \rangle$, $H'_2 = \langle x_1 t^{(1,0)}, x_2 \rangle \leq M'$, and
- $H''_1 = \langle x_3, x_4 \rangle$, $H''_2 = \langle x_3 t^{(0,1)}, x_4 \rangle \leq M''$.

We have $H'_1 \cap H'_2 = \langle x_1^{-i} x_2 x_1^i, i \in \mathbb{Z} \rangle$, $H''_1 \cap H''_2 = \langle x_3^{-i} x_4 x_3^i, i \in \mathbb{Z} \rangle$, and

$\langle H'_1 \cap H'_2, H''_1 \cap H''_2 \rangle = (H'_1 \cap H'_2) * (H''_1 \cap H''_2) = \langle x_1^{-i} x_2 x_1^i, x_3^{-i} x_4 x_3^i \mid i \in \mathbb{Z} \rangle$,

which **does not contain** $x_3^{-1} x_2 x_3 \in \langle H'_1, H''_1 \rangle = \langle x_1, x_2, x_3, x_4 \rangle$
 $\in \langle H'_2, H''_2 \rangle = \langle x_1 t^{(1,0)}, x_2, x_3 t^{(0,1)}, x_4 \rangle$.

Free-times-free-abelian groups

Observation

The same is **not true** in $\mathbb{G} = \mathbb{F}_n \times \mathbb{Z}^m$, even with $M', M'' \leq \mathbb{G}$ in strongly complementary position, i.e., $\langle M'\pi, M''\pi \rangle = M'\pi * M''\pi$ and $\langle M'\tau, M''\tau \rangle = M'\tau \oplus M''\tau$.

Example

Consider $\mathbb{G} = \mathbb{F}_4 \times \mathbb{Z}^2 = \langle x_1, x_2, x_3, x_4 \mid - \rangle \times \langle t_1, t_2 \mid [t_1, t_2] \rangle$,
 $M' = \langle x_1, x_2, t^{(1,0)} \rangle$, $M'' = \langle x_3, x_4, t^{(0,1)} \rangle$, and the respective subgroups

- $H'_1 = \langle x_1, x_2 \rangle$, $H'_2 = \langle x_1 t^{(1,0)}, x_2 \rangle \leq M'$, and
- $H''_1 = \langle x_3, x_4 \rangle$, $H''_2 = \langle x_3 t^{(0,1)}, x_4 \rangle \leq M''$.

We have $H'_1 \cap H'_2 = \langle x_1^{-i} x_2 x_1^i, i \in \mathbb{Z} \rangle$, $H''_1 \cap H''_2 = \langle x_3^{-i} x_4 x_3^i, i \in \mathbb{Z} \rangle$, and

$\langle H'_1 \cap H'_2, H''_1 \cap H''_2 \rangle = (H'_1 \cap H'_2) * (H''_1 \cap H''_2) = \langle x_1^{-i} x_2 x_1^i, x_3^{-i} x_4 x_3^i \mid i \in \mathbb{Z} \rangle$,

which **does not contain** $x_3^{-1} x_2 x_3 \in \langle H'_1, H''_1 \rangle = \langle x_1, x_2, x_3, x_4 \rangle$
 $\in \langle H'_2, H''_2 \rangle = \langle x_1 t^{(1,0)}, x_2, x_3 t^{(0,1)}, x_4 \rangle$.

Free-times-free-abelian groups

Theorem

Let $H'_1, \dots, H'_k \leq G' = \mathbb{F}_{n'} \times \mathbb{Z}^{m'}$ and $H''_1, \dots, H''_k \leq G'' = \mathbb{F}_{n''} \times \mathbb{Z}^{m''}$ be $k \geq 2$ subgroups of G' and G'' , resp. Write $r' = \text{rk}(\bigcap_{j=1}^k H'_j \pi)$, $r'' = \text{rk}(\bigcap_{j=1}^k H''_j \pi)$, and consider $\langle H'_1, H''_1 \rangle, \dots, \langle H'_k, H''_k \rangle \leq G' \circledast G'' = (\mathbb{F}_{n'} \circledast \mathbb{F}_{n''}) \times (\mathbb{Z}^{m'} \oplus \mathbb{Z}^{m''})$. Then, if $\min(r', r'') \neq 1$:

$\bigcap_{j=1}^k \langle H'_j, H''_j \rangle$ is f.g. \Leftrightarrow both $\bigcap_{j=1}^k H'_j$ and $\bigcap_{j=1}^k H''_j$ are f.g.

Observation

Again, *not true* without the hypothesis $\min(r', r'') \neq 1$.

Free-times-free-abelian groups

Theorem

Let $H'_1, \dots, H'_k \leq G' = \mathbb{F}_{n'} \times \mathbb{Z}^{m'}$ and $H''_1, \dots, H''_k \leq G'' = \mathbb{F}_{n''} \times \mathbb{Z}^{m''}$ be $k \geq 2$ subgroups of G' and G'' , resp. Write $r' = \text{rk}(\bigcap_{j=1}^k H'_j \pi)$, $r'' = \text{rk}(\bigcap_{j=1}^k H''_j \pi)$, and consider $\langle H'_1, H''_1 \rangle, \dots, \langle H'_k, H''_k \rangle \leq G' \circledast G'' = (\mathbb{F}_{n'} \circledast \mathbb{F}_{n''}) \times (\mathbb{Z}^{m'} \oplus \mathbb{Z}^{m''})$. Then, if $\min(r', r'') \neq 1$:

$\bigcap_{j=1}^k \langle H'_j, H''_j \rangle$ is f.g. \Leftrightarrow both $\bigcap_{j=1}^k H'_j$ and $\bigcap_{j=1}^k H''_j$ are f.g.

Observation

Again, **not true** without the hypothesis $\min(r', r'') \neq 1$.

Outline

- 1 Our main results
- 2 Free-times-free-abelian groups
- 3 Realizable / unrealizable k -configurations**
- 4 The free case
- 5 Open questions


Positive results

Definition

Define the *join* of two k -configurations χ and χ' as

$$\chi \vee \chi': \mathcal{P}([k]) \setminus \{\emptyset\} \rightarrow \{0, 1\}$$

$$I \mapsto \begin{cases} 0 & \text{if } (I)\chi = (I)\chi' = 0, \\ 1 & \text{otherwise.} \end{cases}$$



Proposition

Let χ' (resp. χ'') be k -config. realized by $H'_1, \dots, H'_k \leq \mathbb{G}' = \mathbb{F}_{n'} \times \mathbb{Z}^{m'}$ (resp. $H''_1, \dots, H''_k \leq \mathbb{G}'' = \mathbb{F}_{n''} \times \mathbb{Z}^{m''}$) with $r'_I = \text{rk}(\bigcap_{i \in I} H'_i \pi) \neq 1$ (resp. $r''_I \neq 1$) $\forall I \subseteq [k]$ with $|I| \geq 2$. Then, $\chi' \vee \chi''$ is realizable in $\mathbb{G}' \otimes \mathbb{G}'' = \mathbb{F}_{n'+n''} \times \mathbb{Z}^{m'+m''}$ by $H_1 = \langle H'_1, H''_1 \rangle, \dots, H_k = \langle H'_k, H''_k \rangle$, again satisfying $r_I \neq 1 \forall I \subseteq [k]$ with $|I| \geq 2$.


Positive results

Definition

Define the *join* of two k -configurations χ and χ' as

$$\chi \vee \chi' : \mathcal{P}([k]) \setminus \{\emptyset\} \rightarrow \{0, 1\}$$

$$I \mapsto \begin{cases} 0 & \text{if } (I)\chi = (I)\chi' = 0, \\ 1 & \text{otherwise.} \end{cases}$$



Proposition

Let χ' (resp. χ'') be k -config. realized by $H'_1, \dots, H'_k \leq \mathbb{G}' = \mathbb{F}_{n'} \times \mathbb{Z}^{m'}$ (resp. $H''_1, \dots, H''_k \leq \mathbb{G}'' = \mathbb{F}_{n''} \times \mathbb{Z}^{m''}$) with $r'_I = \text{rk}(\bigcap_{i \in I} H'_i \pi) \neq 1$ (resp. $r''_I \neq 1$) $\forall I \subseteq [k]$ with $|I| \geq 2$. Then, $\chi' \vee \chi''$ is realizable in $\mathbb{G}' \otimes \mathbb{G}'' = \mathbb{F}_{n'+n''} \times \mathbb{Z}^{m'+m''}$ by $H_1 = \langle H'_1, H''_1 \rangle, \dots, H_k = \langle H'_k, H''_k \rangle$, again satisfying $r_I \neq 1 \forall I \subseteq [k]$ with $|I| \geq 2$.

Positive results

Proposition

The k -config. $\chi_{[k]}$ is realizable in $\mathbb{F}_n \times \mathbb{Z}^{k-1}$.

(Sketch of proof)

$$H_1 = \langle x, y; t^{\mathbf{e}_2}, \dots, t^{\mathbf{e}_{k-1}} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^{k-1},$$

$$H_2 = \langle x, y; t^{\mathbf{e}_1}, t^{\mathbf{e}_3}, \dots, t^{\mathbf{e}_{k-1}} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^{k-1},$$

⋮

$$H_{k-1} = \langle x, y; t^{\mathbf{e}_1}, \dots, t^{\mathbf{e}_{k-2}} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^{k-1},$$

$$H_k = \langle x, yt^{\mathbf{e}_1}; t^{\mathbf{e}_2 - \mathbf{e}_1}, \dots, t^{\mathbf{e}_{k-1} - \mathbf{e}_1} \rangle = \langle x, yt^{\mathbf{e}_1}, \dots, yt^{\mathbf{e}_{k-1}} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^{k-1}$$

Corollary

Any almost-zero k -config. χ_{I_0} is realizable in $\mathbb{F}_n \times \mathbb{Z}^{|I_0|-1}$ by subgroups H_1, \dots, H_k further satisfying $\text{rk}(\bigcap_{i \in I} H_i \pi) \neq 1$, for every $\emptyset \neq I \subseteq [k]$.

Positive results

Proposition

The k -config. $\chi_{[k]}$ is realizable in $\mathbb{F}_n \times \mathbb{Z}^{k-1}$.

(Sketch of proof)

$$H_1 = \langle x, y; t^{\mathbf{e}_2}, \dots, t^{\mathbf{e}_{k-1}} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^{k-1},$$

$$H_2 = \langle x, y; t^{\mathbf{e}_1}, t^{\mathbf{e}_3}, \dots, t^{\mathbf{e}_{k-1}} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^{k-1},$$

$$\vdots$$

$$H_{k-1} = \langle x, y; t^{\mathbf{e}_1}, \dots, t^{\mathbf{e}_{k-2}} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^{k-1},$$

$$H_k = \langle x, yt^{\mathbf{e}_1}; t^{\mathbf{e}_2 - \mathbf{e}_1}, \dots, t^{\mathbf{e}_{k-1} - \mathbf{e}_1} \rangle = \langle x, yt^{\mathbf{e}_1}, \dots, yt^{\mathbf{e}_{k-1}} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^{k-1}.$$

Corollary

Any almost-zero k -config. χ_{I_0} is realizable in $\mathbb{F}_n \times \mathbb{Z}^{|I_0|-1}$ by subgroups H_1, \dots, H_k further satisfying $\text{rk}(\bigcap_{i \in I} H_i \pi) \neq 1$, for every $\emptyset \neq I \subseteq [k]$.

Positive results

Proposition

The k -config. $\chi_{[k]}$ is realizable in $\mathbb{F}_n \times \mathbb{Z}^{k-1}$.

(Sketch of proof)

$$H_1 = \langle x, y; t^{\mathbf{e}_2}, \dots, t^{\mathbf{e}_{k-1}} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^{k-1},$$

$$H_2 = \langle x, y; t^{\mathbf{e}_1}, t^{\mathbf{e}_3}, \dots, t^{\mathbf{e}_{k-1}} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^{k-1},$$

$$\vdots$$

$$H_{k-1} = \langle x, y; t^{\mathbf{e}_1}, \dots, t^{\mathbf{e}_{k-2}} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^{k-1},$$

$$H_k = \langle x, yt^{\mathbf{e}_1}; t^{\mathbf{e}_2 - \mathbf{e}_1}, \dots, t^{\mathbf{e}_{k-1} - \mathbf{e}_1} \rangle = \langle x, yt^{\mathbf{e}_1}, \dots, yt^{\mathbf{e}_{k-1}} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^{k-1}.$$

Corollary

Any almost-zero k -config. χ_{I_0} is realizable in $\mathbb{F}_n \times \mathbb{Z}^{|I_0|-1}$ by subgroups H_1, \dots, H_k further satisfying $\text{rk}(\bigcap_{i \in I} H_i \pi) \neq 1$, for every $\emptyset \neq I \subseteq [k]$.

Positive results

Theorem (Delgado–Roy–V. '22)

Every k -configuration $\chi_{\mathcal{I}}$ is realizable in $\mathbb{F}_n \times \mathbb{Z}^m$, for $n \geq 2$ and $m \geq \sum_{I \in \mathcal{I}} (|I| - 1)$.

(proof)

- Decompose $\chi_{\mathcal{I}} = \chi_{I_1} \vee \cdots \vee \chi_{I_r}$, where $\mathcal{I} = \{I_1, \dots, I_r\}$;
- realize each χ_{I_j} in $\mathbb{F}_2 \times \mathbb{Z}^{|I_j|-1}$, $j = 1, \dots, r$;
- put together in a strongly complementary way. □

Example

Consider $\chi = \chi_{\mathcal{I}}$, where $\mathcal{I} = \{\{1\}, \{2, 3\}, \{1, 3, 4\}, \{2, 3, 4\}\}$. Let us realize it in $\mathbb{F}_2 \times \mathbb{Z}^m$ for $m = 0 + 1 + 2 + 2 = 5$. Decomposing χ , we have

$$\chi = \chi_{\{1\}} \vee \chi_{\{2,3\}} \vee \chi_{\{1,3,4\}} \vee \chi_{\{2,3,4\}}.$$

Positive results

Theorem (Delgado–Roy–V. '22)

Every k -configuration $\chi_{\mathcal{I}}$ is realizable in $\mathbb{F}_n \times \mathbb{Z}^m$, for $n \geq 2$ and $m \geq \sum_{I \in \mathcal{I}} (|I| - 1)$.

(proof)

- Decompose $\chi_{\mathcal{I}} = \chi_{I_1} \vee \cdots \vee \chi_{I_r}$, where $\mathcal{I} = \{I_1, \dots, I_r\}$;
- realize each χ_{I_j} in $\mathbb{F}_2 \times \mathbb{Z}^{|I_j|-1}$, $j = 1, \dots, r$;
- put together in a strongly complementary way. □

Example

Consider $\chi = \chi_{\mathcal{I}}$, where $\mathcal{I} = \{\{1\}, \{2, 3\}, \{1, 3, 4\}, \{2, 3, 4\}\}$. Let us realize it in $\mathbb{F}_2 \times \mathbb{Z}^m$ for $m = 0 + 1 + 2 + 2 = 5$. Decomposing χ , we have

$$\chi = \chi_{\{1\}} \vee \chi_{\{2,3\}} \vee \chi_{\{1,3,4\}} \vee \chi_{\{2,3,4\}}.$$

Positive results

Theorem (Delgado–Roy–V. '22)

Every k -configuration $\chi_{\mathcal{I}}$ is realizable in $\mathbb{F}_n \times \mathbb{Z}^m$, for $n \geq 2$ and $m \geq \sum_{I \in \mathcal{I}} (|I| - 1)$.

(proof)

- Decompose $\chi_{\mathcal{I}} = \chi_{I_1} \vee \cdots \vee \chi_{I_r}$, where $\mathcal{I} = \{I_1, \dots, I_r\}$;
- realize each χ_{I_j} in $\mathbb{F}_2 \times \mathbb{Z}^{|I_j|-1}$, $j = 1, \dots, r$;
- put together in a strongly complementary way. □

Example

Consider $\chi = \chi_{\mathcal{I}}$, where $\mathcal{I} = \{\{1\}, \{2, 3\}, \{1, 3, 4\}, \{2, 3, 4\}\}$. Let us realize it in $\mathbb{F}_2 \times \mathbb{Z}^m$ for $m = 0 + 1 + 2 + 2 = 5$. Decomposing χ , we have

$$\chi = \chi_{\{1\}} \vee \chi_{\{2,3\}} \vee \chi_{\{1,3,4\}} \vee \chi_{\{2,3,4\}}.$$

Positive results

Theorem (Delgado–Roy–V. '22)

Every k -configuration $\chi_{\mathcal{I}}$ is realizable in $\mathbb{F}_n \times \mathbb{Z}^m$, for $n \geq 2$ and $m \geq \sum_{I \in \mathcal{I}} (|I| - 1)$.

(proof)

- Decompose $\chi_{\mathcal{I}} = \chi_{I_1} \vee \cdots \vee \chi_{I_r}$, where $\mathcal{I} = \{I_1, \dots, I_r\}$;
- realize each χ_{I_j} in $\mathbb{F}_2 \times \mathbb{Z}^{|I_j|-1}$, $j = 1, \dots, r$;
- put together in a strongly complementary way. □

Example

Consider $\chi = \chi_{\mathcal{I}}$, where $\mathcal{I} = \{\{1\}, \{2, 3\}, \{1, 3, 4\}, \{2, 3, 4\}\}$. Let us realize it in $\mathbb{F}_2 \times \mathbb{Z}^m$ for $m = 0 + 1 + 2 + 2 = 5$. Decomposing χ , we have

$$\chi = \chi_{\{1\}} \vee \chi_{\{2,3\}} \vee \chi_{\{1,3,4\}} \vee \chi_{\{2,3,4\}}.$$

Positive results

Example (cont.)

In $\mathbb{F}_2 = \langle x, y \mid - \rangle$ take the freely independent words $u_j = y^{-j}xy^j \in \mathbb{F}_2$, $j \in \mathbb{Z}$. Let $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4, \mathbf{e}_5\}$ be the canonical basis for \mathbb{Z}^5 . Realize:

- $\chi_{\{1\}}$ as $H'_1 = \langle \dots, u_{-2}, u_{-1} \rangle$, $H'_2 = \{1\}$, $H'_3 = \{1\}$, $H'_4 = \{1\}$, all inside $G' = \langle \dots, u_{-2}, u_{-1}; - \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
 - $\chi_{\{2,3\}}$ as $H''_1 = \{1\}$, $H''_2 = \langle u_0, u_1 \rangle$, $H''_3 = \langle u_0, u_1 t^{\mathbf{e}_1} \rangle$, $H''_4 = \{1\}$, all inside $G'' = \langle u_0, u_1; t^{\mathbf{e}_1} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
 - $\chi_{\{1,3,4\}}$ as $H'''_1 = \langle u_2, u_3; t^{\mathbf{e}_3} \rangle$, $H'''_2 = \{1\}$, $H'''_3 = \langle u_2, u_3; t^{\mathbf{e}_2} \rangle$, $H'''_4 = \langle u_2, u_3 t^{\mathbf{e}_2}; t^{\mathbf{e}_3 - \mathbf{e}_2} \rangle$, all inside $G''' = \langle u_2, u_3; t^{\mathbf{e}_2}, t^{\mathbf{e}_3} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
 - $\chi_{\{2,3,4\}}$ as $H''''_1 = \{1\}$, $H''''_2 = \langle u_4, u_5; t^{\mathbf{e}_5} \rangle$, $H''''_3 = \langle u_4, u_5; t^{\mathbf{e}_4} \rangle$, $H''''_4 = \langle u_4, u_5 t^{\mathbf{e}_4}; t^{\mathbf{e}_5 - \mathbf{e}_4} \rangle$, all inside $G'''' = \langle u_4, u_5; t^{\mathbf{e}_4}, t^{\mathbf{e}_5} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$.
- And note that $\text{rk}(\bigcap_{i \in I} H'_i \pi) \neq 1$, $\text{rk}(\bigcap_{i \in I} H''_i \pi) \neq 1$, $\text{rk}(\bigcap_{i \in I} H'''_i \pi) \neq 1$, and $\text{rk}(\bigcap_{i \in I} H''''_i \pi) \neq 1$. Therefore, we can realize χ by the following subgroups

Positive results

Example (cont.)

In $\mathbb{F}_2 = \langle x, y \mid - \rangle$ take the freely independent words $u_j = y^{-j}xy^j \in \mathbb{F}_2$, $j \in \mathbb{Z}$. Let $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4, \mathbf{e}_5\}$ be the canonical basis for \mathbb{Z}^5 . Realize:

- $\chi_{\{1\}}$ as $H'_1 = \langle \dots, u_{-2}, u_{-1} \rangle$, $H'_2 = \{1\}$, $H'_3 = \{1\}$, $H'_4 = \{1\}$, all inside $G' = \langle \dots, u_{-2}, u_{-1}; - \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
 - $\chi_{\{2,3\}}$ as $H''_1 = \{1\}$, $H''_2 = \langle u_0, u_1 \rangle$, $H''_3 = \langle u_0, u_1 t^{\mathbf{e}_1} \rangle$, $H''_4 = \{1\}$, all inside $G'' = \langle u_0, u_1; t^{\mathbf{e}_1} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
 - $\chi_{\{1,3,4\}}$ as $H'''_1 = \langle u_2, u_3; t^{\mathbf{e}_3} \rangle$, $H'''_2 = \{1\}$, $H'''_3 = \langle u_2, u_3; t^{\mathbf{e}_2} \rangle$, $H'''_4 = \langle u_2, u_3 t^{\mathbf{e}_2}; t^{\mathbf{e}_3 - \mathbf{e}_2} \rangle$, all inside $G''' = \langle u_2, u_3; t^{\mathbf{e}_2}, t^{\mathbf{e}_3} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
 - $\chi_{\{2,3,4\}}$ as $H''''_1 = \{1\}$, $H''''_2 = \langle u_4, u_5; t^{\mathbf{e}_5} \rangle$, $H''''_3 = \langle u_4, u_5; t^{\mathbf{e}_4} \rangle$, $H''''_4 = \langle u_4, u_5 t^{\mathbf{e}_4}; t^{\mathbf{e}_5 - \mathbf{e}_4} \rangle$, all inside $G'''' = \langle u_4, u_5; t^{\mathbf{e}_4}, t^{\mathbf{e}_5} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$.
- And note that $\text{rk}(\bigcap_{i \in I} H'_i \pi) \neq 1$, $\text{rk}(\bigcap_{i \in I} H''_i \pi) \neq 1$, $\text{rk}(\bigcap_{i \in I} H'''_i \pi) \neq 1$, and $\text{rk}(\bigcap_{i \in I} H''''_i \pi) \neq 1$. Therefore, we can realize χ by the following subgroups

Positive results

Example (cont.)

In $\mathbb{F}_2 = \langle x, y \mid - \rangle$ take the freely independent words $u_j = y^{-j}xy^j \in \mathbb{F}_2$, $j \in \mathbb{Z}$. Let $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4, \mathbf{e}_5\}$ be the canonical basis for \mathbb{Z}^5 . Realize:

- $\chi_{\{1\}}$ as $H'_1 = \langle \dots, u_{-2}, u_{-1} \rangle$, $H'_2 = \{1\}$, $H'_3 = \{1\}$, $H'_4 = \{1\}$, all inside $G' = \langle \dots, u_{-2}, u_{-1}; - \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
 - $\chi_{\{2,3\}}$ as $H''_1 = \{1\}$, $H''_2 = \langle u_0, u_1 \rangle$, $H''_3 = \langle u_0, u_1 t^{\mathbf{e}_1} \rangle$, $H''_4 = \{1\}$, all inside $G'' = \langle u_0, u_1; t^{\mathbf{e}_1} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
 - $\chi_{\{1,3,4\}}$ as $H'''_1 = \langle u_2, u_3; t^{\mathbf{e}_3} \rangle$, $H'''_2 = \{1\}$, $H'''_3 = \langle u_2, u_3; t^{\mathbf{e}_2} \rangle$, $H'''_4 = \langle u_2, u_3 t^{\mathbf{e}_2}; t^{\mathbf{e}_3 - \mathbf{e}_2} \rangle$, all inside $G''' = \langle u_2, u_3; t^{\mathbf{e}_2}, t^{\mathbf{e}_3} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
 - $\chi_{\{2,3,4\}}$ as $H''''_1 = \{1\}$, $H''''_2 = \langle u_4, u_5; t^{\mathbf{e}_5} \rangle$, $H''''_3 = \langle u_4, u_5; t^{\mathbf{e}_4} \rangle$, $H''''_4 = \langle u_4, u_5 t^{\mathbf{e}_4}; t^{\mathbf{e}_5 - \mathbf{e}_4} \rangle$, all inside $G'''' = \langle u_4, u_5; t^{\mathbf{e}_4}, t^{\mathbf{e}_5} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$.
- And note that $\text{rk}(\bigcap_{i \in I} H'_i \pi) \neq 1$, $\text{rk}(\bigcap_{i \in I} H''_i \pi) \neq 1$, $\text{rk}(\bigcap_{i \in I} H'''_i \pi) \neq 1$, and $\text{rk}(\bigcap_{i \in I} H''''_i \pi) \neq 1$. Therefore, we can realize χ by the following subgroups

Positive results

Example (cont.)

In $\mathbb{F}_2 = \langle x, y \mid - \rangle$ take the freely independent words $u_j = y^{-j}xy^j \in \mathbb{F}_2$, $j \in \mathbb{Z}$. Let $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4, \mathbf{e}_5\}$ be the canonical basis for \mathbb{Z}^5 . Realize:

- $\chi_{\{1\}}$ as $H'_1 = \langle \dots, u_{-2}, u_{-1} \rangle$, $H'_2 = \{1\}$, $H'_3 = \{1\}$, $H'_4 = \{1\}$, all inside $G' = \langle \dots, u_{-2}, u_{-1}; - \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
 - $\chi_{\{2,3\}}$ as $H''_1 = \{1\}$, $H''_2 = \langle u_0, u_1 \rangle$, $H''_3 = \langle u_0, u_1 t^{\mathbf{e}_1} \rangle$, $H''_4 = \{1\}$, all inside $G'' = \langle u_0, u_1; t^{\mathbf{e}_1} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
 - $\chi_{\{1,3,4\}}$ as $H'''_1 = \langle u_2, u_3; t^{\mathbf{e}_3} \rangle$, $H'''_2 = \{1\}$, $H'''_3 = \langle u_2, u_3; t^{\mathbf{e}_2} \rangle$, $H'''_4 = \langle u_2, u_3 t^{\mathbf{e}_2}; t^{\mathbf{e}_3 - \mathbf{e}_2} \rangle$, all inside $G''' = \langle u_2, u_3; t^{\mathbf{e}_2}, t^{\mathbf{e}_3} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
 - $\chi_{\{2,3,4\}}$ as $H''''_1 = \{1\}$, $H''''_2 = \langle u_4, u_5; t^{\mathbf{e}_5} \rangle$, $H''''_3 = \langle u_4, u_5; t^{\mathbf{e}_4} \rangle$, $H''''_4 = \langle u_4, u_5 t^{\mathbf{e}_4}; t^{\mathbf{e}_5 - \mathbf{e}_4} \rangle$, all inside $G'''' = \langle u_4, u_5; t^{\mathbf{e}_4}, t^{\mathbf{e}_5} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$.
- And note that $\text{rk}(\bigcap_{i \in I} H'_i \pi) \neq 1$, $\text{rk}(\bigcap_{i \in I} H''_i \pi) \neq 1$, $\text{rk}(\bigcap_{i \in I} H'''_i \pi) \neq 1$, and $\text{rk}(\bigcap_{i \in I} H''''_i \pi) \neq 1$. Therefore, we can realize χ by the following subgroups

Positive results

Example (cont.)

In $\mathbb{F}_2 = \langle x, y \mid - \rangle$ take the freely independent words $u_j = y^{-j}xy^j \in \mathbb{F}_2$, $j \in \mathbb{Z}$. Let $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4, \mathbf{e}_5\}$ be the canonical basis for \mathbb{Z}^5 . Realize:

- $\chi_{\{1\}}$ as $H'_1 = \langle \dots, u_{-2}, u_{-1} \rangle$, $H'_2 = \{1\}$, $H'_3 = \{1\}$, $H'_4 = \{1\}$, all inside $G' = \langle \dots, u_{-2}, u_{-1}; - \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
- $\chi_{\{2,3\}}$ as $H''_1 = \{1\}$, $H''_2 = \langle u_0, u_1 \rangle$, $H''_3 = \langle u_0, u_1 t^{\mathbf{e}_1} \rangle$, $H''_4 = \{1\}$, all inside $G'' = \langle u_0, u_1; t^{\mathbf{e}_1} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
- $\chi_{\{1,3,4\}}$ as $H'''_1 = \langle u_2, u_3; t^{\mathbf{e}_3} \rangle$, $H'''_2 = \{1\}$, $H'''_3 = \langle u_2, u_3; t^{\mathbf{e}_2} \rangle$, $H'''_4 = \langle u_2, u_3 t^{\mathbf{e}_2}; t^{\mathbf{e}_3 - \mathbf{e}_2} \rangle$, all inside $G''' = \langle u_2, u_3; t^{\mathbf{e}_2}, t^{\mathbf{e}_3} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
- $\chi_{\{2,3,4\}}$ as $H''''_1 = \{1\}$, $H''''_2 = \langle u_4, u_5; t^{\mathbf{e}_5} \rangle$, $H''''_3 = \langle u_4, u_5; t^{\mathbf{e}_4} \rangle$, $H''''_4 = \langle u_4, u_5 t^{\mathbf{e}_4}; t^{\mathbf{e}_5 - \mathbf{e}_4} \rangle$, all inside $G'''' = \langle u_4, u_5; t^{\mathbf{e}_4}, t^{\mathbf{e}_5} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$.

And note that $\text{rk}(\bigcap_{i \in I} H'_i \pi) \neq 1$, $\text{rk}(\bigcap_{i \in I} H''_i \pi) \neq 1$,

$\text{rk}(\bigcap_{i \in I} H'''_i \pi) \neq 1$, and $\text{rk}(\bigcap_{i \in I} H''''_i \pi) \neq 1$. Therefore, we can realize χ by the following subgroups

Positive results

Example (cont.)

In $\mathbb{F}_2 = \langle x, y \mid - \rangle$ take the freely independent words $u_j = y^{-j}xy^j \in \mathbb{F}_2$, $j \in \mathbb{Z}$. Let $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4, \mathbf{e}_5\}$ be the canonical basis for \mathbb{Z}^5 . Realize:

- $\chi_{\{1\}}$ as $H'_1 = \langle \dots, u_{-2}, u_{-1} \rangle$, $H'_2 = \{1\}$, $H'_3 = \{1\}$, $H'_4 = \{1\}$, all inside $G' = \langle \dots, u_{-2}, u_{-1}; - \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
- $\chi_{\{2,3\}}$ as $H''_1 = \{1\}$, $H''_2 = \langle u_0, u_1 \rangle$, $H''_3 = \langle u_0, u_1 t^{\mathbf{e}_1} \rangle$, $H''_4 = \{1\}$, all inside $G'' = \langle u_0, u_1; t^{\mathbf{e}_1} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
- $\chi_{\{1,3,4\}}$ as $H'''_1 = \langle u_2, u_3; t^{\mathbf{e}_3} \rangle$, $H'''_2 = \{1\}$, $H'''_3 = \langle u_2, u_3; t^{\mathbf{e}_2} \rangle$, $H'''_4 = \langle u_2, u_3 t^{\mathbf{e}_2}; t^{\mathbf{e}_3 - \mathbf{e}_2} \rangle$, all inside $G''' = \langle u_2, u_3; t^{\mathbf{e}_2}, t^{\mathbf{e}_3} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
- $\chi_{\{2,3,4\}}$ as $H''''_1 = \{1\}$, $H''''_2 = \langle u_4, u_5; t^{\mathbf{e}_5} \rangle$, $H''''_3 = \langle u_4, u_5; t^{\mathbf{e}_4} \rangle$, $H''''_4 = \langle u_4, u_5 t^{\mathbf{e}_4}; t^{\mathbf{e}_5 - \mathbf{e}_4} \rangle$, all inside $G'''' = \langle u_4, u_5; t^{\mathbf{e}_4}, t^{\mathbf{e}_5} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$.

And note that $\text{rk}(\bigcap_{i \in I} H'_i \pi) \neq 1$, $\text{rk}(\bigcap_{i \in I} H''_i \pi) \neq 1$, $\text{rk}(\bigcap_{i \in I} H'''_i \pi) \neq 1$, and $\text{rk}(\bigcap_{i \in I} H''''_i \pi) \neq 1$. Therefore, we can realize χ by the following subgroups

Positive results

Example (cont.)

In $\mathbb{F}_2 = \langle x, y \mid - \rangle$ take the freely independent words $u_j = y^{-j}xy^j \in \mathbb{F}_2$, $j \in \mathbb{Z}$. Let $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4, \mathbf{e}_5\}$ be the canonical basis for \mathbb{Z}^5 . Realize:

- $\chi_{\{1\}}$ as $H'_1 = \langle \dots, u_{-2}, u_{-1} \rangle$, $H'_2 = \{1\}$, $H'_3 = \{1\}$, $H'_4 = \{1\}$, all inside $G' = \langle \dots, u_{-2}, u_{-1}; - \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
- $\chi_{\{2,3\}}$ as $H''_1 = \{1\}$, $H''_2 = \langle u_0, u_1 \rangle$, $H''_3 = \langle u_0, u_1 t^{\mathbf{e}_1} \rangle$, $H''_4 = \{1\}$, all inside $G'' = \langle u_0, u_1; t^{\mathbf{e}_1} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
- $\chi_{\{1,3,4\}}$ as $H'''_1 = \langle u_2, u_3; t^{\mathbf{e}_3} \rangle$, $H'''_2 = \{1\}$, $H'''_3 = \langle u_2, u_3; t^{\mathbf{e}_2} \rangle$, $H'''_4 = \langle u_2, u_3 t^{\mathbf{e}_2}; t^{\mathbf{e}_3 - \mathbf{e}_2} \rangle$, all inside $G''' = \langle u_2, u_3; t^{\mathbf{e}_2}, t^{\mathbf{e}_3} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$;
- $\chi_{\{2,3,4\}}$ as $H''''_1 = \{1\}$, $H''''_2 = \langle u_4, u_5; t^{\mathbf{e}_5} \rangle$, $H''''_3 = \langle u_4, u_5; t^{\mathbf{e}_4} \rangle$, $H''''_4 = \langle u_4, u_5 t^{\mathbf{e}_4}; t^{\mathbf{e}_5 - \mathbf{e}_4} \rangle$, all inside $G'''' = \langle u_4, u_5; t^{\mathbf{e}_4}, t^{\mathbf{e}_5} \rangle \leq \mathbb{F}_2 \times \mathbb{Z}^5$.

And note that $\text{rk}(\bigcap_{i \in I} H'_i \pi) \neq 1$, $\text{rk}(\bigcap_{i \in I} H''_i \pi) \neq 1$, $\text{rk}(\bigcap_{i \in I} H'''_i \pi) \neq 1$, and $\text{rk}(\bigcap_{i \in I} H''''_i \pi) \neq 1$. Therefore, we can realize χ by the following subgroups

Positive results

Example (cont.)

$$H_1 = \langle \dots, u_{-2}, u_{-1}, u_2, u_3; t^{\mathbf{e}_3} \rangle,$$

$$H_2 = \langle u_0, u_1, u_4, u_5; t^{\mathbf{e}_5} \rangle,$$

$$H_3 = \langle u_0, u_1 t^{\mathbf{e}_1}, u_2, u_3, u_4, u_5; t^{\mathbf{e}_2}, t^{\mathbf{e}_4} \rangle,$$

$$H_4 = \langle u_2, u_3 t^{\mathbf{e}_2}, u_4, u_5 t^{\mathbf{e}_4}; t^{\mathbf{e}_3 - \mathbf{e}_2}, t^{\mathbf{e}_5 - \mathbf{e}_4} \rangle.$$

of $G' * G'' * G''' * G'''' \leq \mathbb{F}_2 \times \mathbb{Z}^5$. □

Corollary

$\mathbb{F}_2 \times (\bigoplus_{\mathbb{N}_0} \mathbb{Z})$ is intersection-saturated.

Theorem (Delgado–Roy–V. '22)

There exist finitely presented intersection-saturated groups.

Positive results

Example (cont.)

$$H_1 = \langle \dots, u_{-2}, u_{-1}, u_2, u_3; t^{\mathbf{e}_3} \rangle,$$

$$H_2 = \langle u_0, u_1, u_4, u_5; t^{\mathbf{e}_5} \rangle,$$

$$H_3 = \langle u_0, u_1 t^{\mathbf{e}_1}, u_2, u_3, u_4, u_5; t^{\mathbf{e}_2}, t^{\mathbf{e}_4} \rangle,$$

$$H_4 = \langle u_2, u_3 t^{\mathbf{e}_2}, u_4, u_5 t^{\mathbf{e}_4}; t^{\mathbf{e}_3 - \mathbf{e}_2}, t^{\mathbf{e}_5 - \mathbf{e}_4} \rangle.$$

of $G' \ast G'' \ast G''' \ast G'''' \leq \mathbb{F}_2 \times \mathbb{Z}^5$. □

Corollary

$\mathbb{F}_2 \times \left(\bigoplus_{\mathbb{N}_0} \mathbb{Z} \right)$ is intersection-saturated.

Theorem (Delgado–Roy–V. '22)

There exist finitely presented intersection-saturated groups.

Positive results

Example (cont.)

$$H_1 = \langle \dots, u_{-2}, u_{-1}, u_2, u_3; t^{\mathbf{e}_3} \rangle,$$

$$H_2 = \langle u_0, u_1, u_4, u_5; t^{\mathbf{e}_5} \rangle,$$

$$H_3 = \langle u_0, u_1 t^{\mathbf{e}_1}, u_2, u_3, u_4, u_5; t^{\mathbf{e}_2}, t^{\mathbf{e}_4} \rangle,$$

$$H_4 = \langle u_2, u_3 t^{\mathbf{e}_2}, u_4, u_5 t^{\mathbf{e}_4}; t^{\mathbf{e}_3 - \mathbf{e}_2}, t^{\mathbf{e}_5 - \mathbf{e}_4} \rangle.$$

of $G' \ast G'' \ast G''' \ast G'''' \leq \mathbb{F}_2 \times \mathbb{Z}^5$. □

Corollary

$\mathbb{F}_2 \times \left(\bigoplus_{\mathbb{N}_0} \mathbb{Z} \right)$ is intersection-saturated.

Theorem (Delgado–Roy–V. '22)

There exist finitely presented intersection-saturated groups.

Positive results

Theorem (Delgado–Roy–V. '22)

There exist finitely presented intersection-saturated groups.

(Proof 1)

- Consider Thomson's group F ;
- it is well known to be finitely presented and to contain $\bigoplus_{\mathbb{N}_0} \mathbb{Z}$;
- therefore, $\mathbb{F}_2 \times F$ is intersection-saturated. □
- (Need to take $\mathbb{F}_2 \times$ because F does not contain \mathbb{F}_2 .)

(Proof 2)

- Consider $G = \left(\bigoplus_{\mathbb{N}_0} \mathbb{Z} \right) \rtimes_{\alpha} \mathbb{Z}$, where α is the automorphism given by right translation of generators;
- G is recursively presented so, it embeds in a finitely presented group, $G \hookrightarrow G'$;
- $\mathbb{F}_2 \times G'$ is finitely presented and intersection-saturated. □

Positive results

Theorem (Delgado–Roy–V. '22)

There exist finitely presented intersection-saturated groups.

(Proof 1)

- *Consider Thomson's group F ;*
- *it is well known to be finitely presented and to contain $\bigoplus_{\mathbb{N}_0} \mathbb{Z}$;*
- *therefore, $\mathbb{F}_2 \times F$ is intersection-saturated. □*
- *(Need to take $\mathbb{F}_2 \times$ because F does not contain \mathbb{F}_2 .)*

(Proof 2)

- *Consider $G = \left(\bigoplus_{\mathbb{N}_0} \mathbb{Z} \right) \rtimes_{\alpha} \mathbb{Z}$, where α is the automorphism given by right translation of generators;*
- *G is recursively presented so, it embeds in a finitely presented group, $G \hookrightarrow G'$;*
- *$\mathbb{F}_2 \times G'$ is finitely presented and intersection-saturated. □*

Positive results

Theorem (Delgado–Roy–V. '22)

There exist finitely presented intersection-saturated groups.

(Proof 1)

- Consider Thomson's group F ;
- it is well known to be finitely presented and to contain $\bigoplus_{\mathbb{N}_0} \mathbb{Z}$;
- therefore, $\mathbb{F}_2 \times F$ is intersection-saturated.
- (Need to take $\mathbb{F}_2 \times$ because F does not contain \mathbb{F}_2 .)

(Proof 2)

- Consider $G = \left(\bigoplus_{\mathbb{N}_0} \mathbb{Z} \right) \rtimes_{\alpha} \mathbb{Z}$, where α is the automorphism given by right translation of generators;
- G is recursively presented so, it embeds in a finitely presented group, $G \hookrightarrow G'$;
- $\mathbb{F}_2 \times G'$ is finitely presented and intersection-saturated.

Positive results

Theorem (Delgado–Roy–V. '22)

There exist finitely presented intersection-saturated groups.

(Proof 1)

- Consider Thomson's group F ;
- it is well known to be finitely presented and to contain $\bigoplus_{\mathbb{N}_0} \mathbb{Z}$;
- therefore, $\mathbb{F}_2 \times F$ is intersection-saturated. □
- (Need to take $\mathbb{F}_2 \times$ because F does not contain \mathbb{F}_2 .)

(Proof 2)

- Consider $G = \left(\bigoplus_{\mathbb{N}_0} \mathbb{Z} \right) \rtimes_{\alpha} \mathbb{Z}$, where α is the automorphism given by right translation of generators;
- G is recursively presented so, it embeds in a finitely presented group, $G \hookrightarrow G'$;
- $\mathbb{F}_2 \times G'$ is finitely presented and intersection-saturated. □

Positive results

Theorem (Delgado–Roy–V. '22)

There exist finitely presented intersection-saturated groups.

(Proof 1)

- Consider Thomson's group F ;
- it is well known to be finitely presented and to contain $\bigoplus_{\mathbb{N}_0} \mathbb{Z}$;
- therefore, $\mathbb{F}_2 \times F$ is intersection-saturated. □
- (Need to take $\mathbb{F}_2 \times$ because F does not contain \mathbb{F}_2 .)

(Proof 2)

- Consider $G = \left(\bigoplus_{\mathbb{N}_0} \mathbb{Z} \right) \rtimes_{\alpha} \mathbb{Z}$, where α is the automorphism given by right translation of generators;
- G is recursively presented so, it embeds in a finitely presented group, $G \hookrightarrow G'$;
- $\mathbb{F}_2 \times G'$ is finitely presented and intersection-saturated. □

Positive results

Theorem (Delgado–Roy–V. '22)

There exist finitely presented intersection-saturated groups.

(Proof 1)

- Consider Thomson's group F ;
- it is well known to be finitely presented and to contain $\bigoplus_{\mathbb{N}_0} \mathbb{Z}$;
- therefore, $\mathbb{F}_2 \times F$ is intersection-saturated. □
- (Need to take $\mathbb{F}_2 \times$ because F does not contain \mathbb{F}_2 .)

(Proof 2)

- Consider $G = \left(\bigoplus_{\mathbb{N}_0} \mathbb{Z} \right) \rtimes_{\alpha} \mathbb{Z}$, where α is the automorphism given by right translation of generators;
- G is recursively presented so, it embeds in a finitely presented group, $G \hookrightarrow G'$;
- $\mathbb{F}_2 \times G'$ is finitely presented and intersection-saturated. □

Positive results

Theorem (Delgado–Roy–V. '22)

There exist finitely presented intersection-saturated groups.

(Proof 1)

- Consider Thomson's group F ;
- it is well known to be finitely presented and to contain $\bigoplus_{\mathbb{N}_0} \mathbb{Z}$;
- therefore, $\mathbb{F}_2 \times F$ is intersection-saturated. □
- (Need to take $\mathbb{F}_2 \times$ because F does not contain \mathbb{F}_2 .)

(Proof 2)

- Consider $G = \left(\bigoplus_{\mathbb{N}_0} \mathbb{Z} \right) \rtimes_{\alpha} \mathbb{Z}$, where α is the automorphism given by right translation of generators;
- G is recursively presented so, it embeds in a finitely presented group, $G \hookrightarrow G'$;
- $\mathbb{F}_2 \times G'$ is finitely presented and intersection-saturated. □

Positive results

Theorem (Delgado–Roy–V. '22)

There exist finitely presented intersection-saturated groups.

(Proof 1)

- Consider Thomson's group F ;
- it is well known to be finitely presented and to contain $\bigoplus_{\mathbb{N}_0} \mathbb{Z}$;
- therefore, $\mathbb{F}_2 \times F$ is intersection-saturated. □
- (Need to take $\mathbb{F}_2 \times$ because F does not contain \mathbb{F}_2 .)

(Proof 2)

- Consider $G = \left(\bigoplus_{\mathbb{N}_0} \mathbb{Z} \right) \rtimes_{\alpha} \mathbb{Z}$, where α is the automorphism given by right translation of generators;
- G is recursively presented so, it embeds in a finitely presented group, $G \hookrightarrow G'$;
- $\mathbb{F}_2 \times G'$ is finitely presented and intersection-saturated. □

An obstruction


Lemma

Let $H_1, \dots, H_k \leq \mathbb{G} = \mathbb{F}_n \times \mathbb{Z}^m$. Suppose that, for $\emptyset \neq I, J \subseteq [k]$, H_I and H_J are f.g. whereas $H_{I \cup J} = H_I \cap H_J$ is not. Then, $\exists i \in I, \exists j \in J$ s.t. $L_i = H_i \cap \mathbb{Z}^m$ and $L_j = H_j \cap \mathbb{Z}^m$ both have rank strictly smaller than m .

Proposition

Let χ be a k -config. and $\emptyset \neq I_1, \dots, I_r \subseteq [k]$ be $r \geq 2$ subsets s.t. $\forall j \in [r], (I_1 \cup \dots \cup \widehat{I_j} \cup \dots \cup I_r)\chi = \mathbf{0}$, but $(I_1 \cup \dots \cup I_r)\chi = \mathbf{1}$. Then χ is *not realizable* in $\mathbb{F}_n \times \mathbb{Z}^{r-2}$.

Corollary



The 3-configurations

are *not realizable* in $\mathbb{F}_n \times \mathbb{Z}$.

An obstruction


Lemma

Let $H_1, \dots, H_k \leq \mathbb{G} = \mathbb{F}_n \times \mathbb{Z}^m$. Suppose that, for $\emptyset \neq I, J \subseteq [k]$, H_I and H_J are f.g. whereas $H_{I \cup J} = H_I \cap H_J$ is not. Then, $\exists i \in I, \exists j \in J$ s.t. $L_i = H_i \cap \mathbb{Z}^m$ and $L_j = H_j \cap \mathbb{Z}^m$ both have rank strictly smaller than m .

Proposition

Let χ be a k -config. and $\emptyset \neq I_1, \dots, I_r \subseteq [k]$ be $r \geq 2$ subsets s.t. $\forall j \in [r], (I_1 \cup \dots \cup \widehat{I_j} \cup \dots \cup I_r)\chi = \mathbf{0}$, but $(I_1 \cup \dots \cup I_r)\chi = \mathbf{1}$. Then χ is **not realizable** in $\mathbb{F}_n \times \mathbb{Z}^{r-2}$.

Corollary



The 3-configurations

are not realizable in $\mathbb{F}_n \times \mathbb{Z}$.

An obstruction


Lemma

Let $H_1, \dots, H_k \leq \mathbb{G} = \mathbb{F}_n \times \mathbb{Z}^m$. Suppose that, for $\emptyset \neq I, J \subseteq [k]$, H_I and H_J are f.g. whereas $H_{I \cup J} = H_I \cap H_J$ is not. Then, $\exists i \in I, \exists j \in J$ s.t. $L_i = H_i \cap \mathbb{Z}^m$ and $L_j = H_j \cap \mathbb{Z}^m$ both have rank strictly smaller than m .

Proposition

Let χ be a k -config. and $\emptyset \neq I_1, \dots, I_r \subseteq [k]$ be $r \geq 2$ subsets s.t. $\forall j \in [r], (I_1 \cup \dots \cup \widehat{I_j} \cup \dots \cup I_r)\chi = \mathbf{0}$, but $(I_1 \cup \dots \cup I_r)\chi = \mathbf{1}$. Then χ is **not realizable** in $\mathbb{F}_n \times \mathbb{Z}^{r-2}$.

Corollary



The 3-configurations

are **not realizable** in $\mathbb{F}_n \times \mathbb{Z}$.

An obstruction

Proposition

The k -configuration $\chi_{[k]}$ is realizable in $\mathbb{F}_n \times \mathbb{Z}^{k-1}$, but **not** in $\mathbb{F}_n \times \mathbb{Z}^{k-2}$.

Hence, the set of configurations realizable in $\mathbb{F}_n \times \mathbb{Z}^m$ increases strictly with m .

Outline

- 1 Our main results
- 2 Free-times-free-abelian groups
- 3 Realizable / unrealizable k -configurations
- 4 The free case**
- 5 Open questions

More on configurations

Definition

Let χ be a k -config. and let $i \in [k]$. Its *restriction to $\hat{i} = [k] \setminus \{i\}$* is the $(k-1)$ -configuration

$$\begin{aligned} \chi_{|\hat{i}}: \mathcal{P}([k] \setminus \{i\}) \setminus \{\emptyset\} &\rightarrow \{0, 1\} \\ I &\mapsto (I)\chi. \end{aligned}$$

Definition

Given two k -configurations χ, χ' and $\delta \in \{0, 1\}$, we define

$$\begin{aligned} \chi \boxplus_{\delta} \chi': \mathcal{P}([k+1]) \setminus \{\emptyset\} &\rightarrow \{0, 1\} \\ I &\mapsto \begin{cases} (I)\chi & \text{if } k+1 \notin I, \\ (I \setminus \{k+1\})\chi' & \text{if } \{k+1\} \subsetneq I, \\ \delta & \text{if } \{k+1\} = I, \end{cases} \end{aligned}$$

a $(k+1)$ -configuration.

More on configurations

Definition

Let χ be a k -config. and let $i \in [k]$. Its *restriction to $\hat{i} = [k] \setminus \{i\}$* is the $(k-1)$ -configuration

$$\begin{aligned} \chi_{\hat{i}}: \mathcal{P}([k] \setminus \{i\}) \setminus \{\emptyset\} &\rightarrow \{0, 1\} \\ I &\mapsto (I)\chi. \end{aligned}$$

Definition


Given two k -configurations χ, χ' and $\delta \in \{0, 1\}$, we define

$$\begin{aligned} \chi \boxplus_{\delta} \chi': \mathcal{P}([k+1]) \setminus \{\emptyset\} &\rightarrow \{0, 1\} \\ I &\mapsto \begin{cases} (I)\chi & \text{if } k+1 \notin I, \\ (I \setminus \{k+1\})\chi' & \text{if } \{k+1\} \subsetneq I, \\ \delta & \text{if } \{k+1\} = I, \end{cases} \end{aligned}$$

a $(k+1)$ -configuration.

More on configurations

Example



Definition


Let χ be a k -configuration, and $i \in [k]$. The index i is said to be *0-monochromatic (in χ)* if $(I)\chi = 0 \forall I \subseteq [k]$ containing i ; i.e., if $\chi = \chi_{\widehat{i}} \boxplus_0 \mathbf{0}$. Similarly, the index i is said to be *1-monochromatic (in χ)* if $\chi = \chi_{\widehat{i}} \boxplus_1 \mathbf{1}$.

Lemma

If a k -configuration χ is realizable in \mathbb{F}_n with $n \geq 2$, then the $(k+1)$ -configurations $\chi \boxplus_0 \mathbf{0}$, $\chi \boxplus_1 \mathbf{1}$, $\chi \boxplus_0 \chi$, and $\chi \boxplus_1 \chi$ are also realizable in \mathbb{F}_n .

More on configurations

Example



Definition


Let χ be a k -configuration, and $i \in [k]$. The index i is said to be **0-monochromatic** (in χ) if $(I)\chi = 0 \forall I \subseteq [k]$ containing i ; i.e., if $\chi = \chi_{\uparrow \hat{i}} \boxplus_0 \mathbf{0}$. Similarly, the index i is said to be **1-monochromatic** (in χ) if $\chi = \chi_{\uparrow \hat{i}} \boxplus_1 \mathbf{1}$.

Lemma

If a k -configuration χ is realizable in \mathbb{F}_n with $n \geq 2$, then the $(k+1)$ -configurations $\chi \boxplus_0 \mathbf{0}$, $\chi \boxplus_1 \mathbf{1}$, $\chi \boxplus_0 \chi$, and $\chi \boxplus_1 \chi$ are also realizable in \mathbb{F}_n .

More on configurations

Example



Definition

Let χ be a k -configuration, and $i \in [k]$. The index i is said to be **0-monochromatic** (in χ) if $(I)\chi = 0 \forall I \subseteq [k]$ containing i ; i.e., if $\chi = \chi_{\uparrow I} \boxplus_0 \mathbf{0}$. Similarly, the index i is said to be **1-monochromatic** (in χ) if $\chi = \chi_{\uparrow I} \boxplus_1 \mathbf{1}$.

Lemma

If a k -configuration χ is realizable in \mathbb{F}_n with $n \geq 2$, then the $(k+1)$ -configurations $\chi \boxplus_0 \mathbf{0}$, $\chi \boxplus_1 \mathbf{1}$, $\chi \boxplus_0 \chi$, and $\chi \boxplus_1 \chi$ are also realizable in \mathbb{F}_n .

Characterization for the free case

(Proof)

Let $\mathbb{F}_2 * \mathbb{F}_{\mathbb{N}_0} \simeq W * U = \langle w_1, w_2, \dots \rangle * \langle u, v \rangle \leq \mathbb{F}_n$, and take $H_1, \dots, H_k \leq W \leq \mathbb{F}_n$ realizing χ . Now, in order to realize:

- $\chi \boxplus_0 \mathbf{0}$, take $\tilde{H}_1 = H_1, \dots, \tilde{H}_k = H_k$, and $\tilde{H}_{k+1} = \{1\}$;
- $\chi \boxplus_1 \mathbf{1}$, take $\tilde{H}_1 = H_1 * \langle u, v \rangle, \dots, \tilde{H}_k = H_k * \langle u, v \rangle$ and $\tilde{H}_{k+1} = \ll u \gg_U: \tilde{H}_1, \dots, \tilde{H}_k$ realize $\chi \vee \mathbf{0} = \chi$ and, for every $i \neq k+1$, $\tilde{H}_{k+1} \cap \tilde{H}_i = \tilde{H}_{k+1}$ which is non-f.g.;
- $\chi \boxplus_0 \chi$, take $\tilde{H}_1 = H_1, \dots, \tilde{H}_k = H_k$, and $\tilde{H}_{k+1} = \mathbb{F}_n$;
- $\chi \boxplus_1 \chi$, take $\tilde{H}_1 = H_1, \dots, \tilde{H}_k = H_k$, and $\tilde{H}_{k+1} = W$. □

Definition

A k -configuration χ is said to be *Howson* if, for every $\emptyset \neq I, J \subseteq [k]$, $(I)\chi = (J)\chi = \mathbf{0} \Rightarrow (I \cup J)\chi = \mathbf{0}$.

Characterization for the free case

(Proof)

Let $\mathbb{F}_2 * \mathbb{F}_{\mathbb{N}_0} \simeq W * U = \langle w_1, w_2, \dots \rangle * \langle u, v \rangle \leq \mathbb{F}_n$, and take $H_1, \dots, H_k \leq W \leq \mathbb{F}_n$ realizing χ . Now, in order to realize:

- $\chi \boxplus_0 \mathbf{0}$, take $\tilde{H}_1 = H_1, \dots, \tilde{H}_k = H_k$, and $\tilde{H}_{k+1} = \{1\}$;
- $\chi \boxplus_1 \mathbf{1}$, take $\tilde{H}_1 = H_1 * \langle u, v \rangle, \dots, \tilde{H}_k = H_k * \langle u, v \rangle$ and $\tilde{H}_{k+1} = \ll u \gg_U: \tilde{H}_1, \dots, \tilde{H}_k$ realize $\chi \vee \mathbf{0} = \chi$ and, for every $i \neq k+1$, $\tilde{H}_{k+1} \cap \tilde{H}_i = \tilde{H}_{k+1}$ which is non-f.g.;
- $\chi \boxplus_0 \chi$, take $\tilde{H}_1 = H_1, \dots, \tilde{H}_k = H_k$, and $\tilde{H}_{k+1} = \mathbb{F}_n$;
- $\chi \boxplus_1 \chi$, take $\tilde{H}_1 = H_1, \dots, \tilde{H}_k = H_k$, and $\tilde{H}_{k+1} = W$. □

Definition

A k -configuration χ is said to be *Howson* if, for every $\emptyset \neq I, J \subseteq [k]$, $(I)\chi = (J)\chi = \mathbf{0} \Rightarrow (I \cup J)\chi = \mathbf{0}$.

Characterization for the free case

(Proof)

Let $\mathbb{F}_2 * \mathbb{F}_{\mathbb{N}_0} \simeq W * U = \langle w_1, w_2, \dots \rangle * \langle u, v \rangle \leq \mathbb{F}_n$, and take $H_1, \dots, H_k \leq W \leq \mathbb{F}_n$ realizing χ . Now, in order to realize:

- $\chi \boxplus_0 \mathbf{0}$, take $\tilde{H}_1 = H_1, \dots, \tilde{H}_k = H_k$, and $\tilde{H}_{k+1} = \{1\}$;
- $\chi \boxplus_1 \mathbf{1}$, take $\tilde{H}_1 = H_1 * \langle u, v \rangle, \dots, \tilde{H}_k = H_k * \langle u, v \rangle$ and $\tilde{H}_{k+1} = \ll u \gg_U: \tilde{H}_1, \dots, \tilde{H}_k$ realize $\chi \vee \mathbf{0} = \chi$ and, for every $i \neq k+1$, $\tilde{H}_{k+1} \cap \tilde{H}_i = \tilde{H}_{k+1}$ which is non-f.g.;
- $\chi \boxplus_0 \chi$, take $\tilde{H}_1 = H_1, \dots, \tilde{H}_k = H_k$, and $\tilde{H}_{k+1} = \mathbb{F}_n$;
- $\chi \boxplus_1 \chi$, take $\tilde{H}_1 = H_1, \dots, \tilde{H}_k = H_k$, and $\tilde{H}_{k+1} = W$. □

Definition

A k -configuration χ is said to be *Howson* if, for every $\emptyset \neq I, J \subseteq [k]$, $(I)\chi = (J)\chi = \mathbf{0} \Rightarrow (I \cup J)\chi = \mathbf{0}$.

Characterization for the free case

(Proof)

Let $\mathbb{F}_2 * \mathbb{F}_{\mathbb{N}_0} \simeq W * U = \langle w_1, w_2, \dots \rangle * \langle u, v \rangle \leq \mathbb{F}_n$, and take $H_1, \dots, H_k \leq W \leq \mathbb{F}_n$ realizing χ . Now, in order to realize:

- $\chi \boxplus_0 \mathbf{0}$, take $\tilde{H}_1 = H_1, \dots, \tilde{H}_k = H_k$, and $\tilde{H}_{k+1} = \{1\}$;
- $\chi \boxplus_1 \mathbf{1}$, take $\tilde{H}_1 = H_1 * \langle u, v \rangle, \dots, \tilde{H}_k = H_k * \langle u, v \rangle$ and $\tilde{H}_{k+1} = \ll u \gg_U: \tilde{H}_1, \dots, \tilde{H}_k$ realize $\chi \vee \mathbf{0} = \chi$ and, for every $i \neq k+1$, $\tilde{H}_{k+1} \cap \tilde{H}_i = \tilde{H}_{k+1}$ which is non-f.g.;
- $\chi \boxplus_0 \chi$, take $\tilde{H}_1 = H_1, \dots, \tilde{H}_k = H_k$, and $\tilde{H}_{k+1} = \mathbb{F}_n$;
- $\chi \boxplus_1 \chi$, take $\tilde{H}_1 = H_1, \dots, \tilde{H}_k = H_k$, and $\tilde{H}_{k+1} = W$. □

Definition

A k -configuration χ is said to be *Howson* if, for every $\emptyset \neq I, J \subseteq [k]$, $(I)\chi = (J)\chi = \mathbf{0} \Rightarrow (I \cup J)\chi = \mathbf{0}$.

Characterization for the free case

(Proof)

Let $\mathbb{F}_2 * \mathbb{F}_{\mathbb{N}_0} \simeq W * U = \langle w_1, w_2, \dots \rangle * \langle u, v \rangle \leq \mathbb{F}_n$, and take $H_1, \dots, H_k \leq W \leq \mathbb{F}_n$ realizing χ . Now, in order to realize:

- $\chi \boxplus_0 \mathbf{0}$, take $\tilde{H}_1 = H_1, \dots, \tilde{H}_k = H_k$, and $\tilde{H}_{k+1} = \{1\}$;
- $\chi \boxplus_1 \mathbf{1}$, take $\tilde{H}_1 = H_1 * \langle u, v \rangle, \dots, \tilde{H}_k = H_k * \langle u, v \rangle$ and $\tilde{H}_{k+1} = \ll u \gg_U: \tilde{H}_1, \dots, \tilde{H}_k$ realize $\chi \vee \mathbf{0} = \chi$ and, for every $i \neq k+1$, $\tilde{H}_{k+1} \cap \tilde{H}_i = \tilde{H}_{k+1}$ which is non-f.g.;
- $\chi \boxplus_0 \chi$, take $\tilde{H}_1 = H_1, \dots, \tilde{H}_k = H_k$, and $\tilde{H}_{k+1} = \mathbb{F}_n$;
- $\chi \boxplus_1 \chi$, take $\tilde{H}_1 = H_1, \dots, \tilde{H}_k = H_k$, and $\tilde{H}_{k+1} = W$. □

Definition

A k -configuration χ is said to be *Howson* if, for every $\emptyset \neq I, J \subseteq [k]$, $(I)\chi = (J)\chi = \mathbf{0} \Rightarrow (I \cup J)\chi = \mathbf{0}$.

Characterization for the free case

(Proof)

Let $\mathbb{F}_2 * \mathbb{F}_{\mathbb{N}_0} \simeq W * U = \langle w_1, w_2, \dots \rangle * \langle u, v \rangle \leq \mathbb{F}_n$, and take $H_1, \dots, H_k \leq W \leq \mathbb{F}_n$ realizing χ . Now, in order to realize:

- $\chi \boxplus_0 \mathbf{0}$, take $\tilde{H}_1 = H_1, \dots, \tilde{H}_k = H_k$, and $\tilde{H}_{k+1} = \{1\}$;
- $\chi \boxplus_1 \mathbf{1}$, take $\tilde{H}_1 = H_1 * \langle u, v \rangle, \dots, \tilde{H}_k = H_k * \langle u, v \rangle$ and $\tilde{H}_{k+1} = \ll u \gg_U: \tilde{H}_1, \dots, \tilde{H}_k$ realize $\chi \vee \mathbf{0} = \chi$ and, for every $i \neq k+1$, $\tilde{H}_{k+1} \cap \tilde{H}_i = \tilde{H}_{k+1}$ which is non-f.g.;
- $\chi \boxplus_0 \chi$, take $\tilde{H}_1 = H_1, \dots, \tilde{H}_k = H_k$, and $\tilde{H}_{k+1} = \mathbb{F}_n$;
- $\chi \boxplus_1 \chi$, take $\tilde{H}_1 = H_1, \dots, \tilde{H}_k = H_k$, and $\tilde{H}_{k+1} = W$. □

Definition

A k -configuration χ is said to be **Howson** if, for every $\emptyset \neq I, J \subseteq [k]$, $(I)\chi = (J)\chi = \mathbf{0} \Rightarrow (I \cup J)\chi = \mathbf{0}$.

Characterization for the free case

Theorem (Delgado–Roy–V., '22)

A k -configuration is realizable in \mathbb{F}_n , $n \geq 2 \Leftrightarrow$ it is Howson.

(Proof)

For \Leftarrow , we will do induction on the cardinal of the support of χ , say s (regardless of its size k).

- If $s = 0$ then $\chi = \mathbf{0}$, clearly realizable in \mathbb{F}_2 .*
- Given χ with $|\text{supp}(\chi)| = s$ and being Howson, define the **cone** of χ with vertex $I \subseteq [k]$, denoted by $c_I(\chi)$, as*

$$c_I(\chi): \mathcal{P}([k]) \setminus \{\emptyset\} \rightarrow \{0, 1\}$$

$$J \mapsto \begin{cases} 0 & \text{if } J \not\subseteq I, \\ (J)\chi & \text{if } J \subseteq I. \end{cases}$$

- Now let $I_1, \dots, I_p \subseteq [k]$ be the maximal elements in $\text{supp}(\chi)$ (w.r.t. inclusion). It is clear that $\chi = c_{I_1}(\chi) \vee \dots \vee c_{I_p}(\chi)$.*

Characterization for the free case

Theorem (Delgado–Roy–V., '22)

A k -configuration is realizable in \mathbb{F}_n , $n \geq 2 \Leftrightarrow$ it is Howson.

(Proof)

For \Leftarrow , we will do induction on the cardinal of the support of χ , say s (regardless of its size k).

- If $s = 0$ then $\chi = \mathbf{0}$, clearly realizable in \mathbb{F}_2 .*
- Given χ with $|\text{supp}(\chi)| = s$ and being Howson, define the *cone* of χ with vertex $I \subseteq [k]$, denoted by $c_I(\chi)$, as*

$$c_I(\chi): \mathcal{P}([k]) \setminus \{\emptyset\} \rightarrow \{0, 1\}$$

$$J \mapsto \begin{cases} 0 & \text{if } J \not\subseteq I, \\ (J)\chi & \text{if } J \subseteq I. \end{cases}$$

- Now let $I_1, \dots, I_p \subseteq [k]$ be the maximal elements in $\text{supp}(\chi)$ (w.r.t. inclusion). It is clear that $\chi = c_{I_1}(\chi) \vee \dots \vee c_{I_p}(\chi)$.*

Characterization for the free case

Theorem (Delgado–Roy–V., '22)

A k -configuration is realizable in \mathbb{F}_n , $n \geq 2 \Leftrightarrow$ it is Howson.

(Proof)

For \Leftarrow , we will do induction on the cardinal of the support of χ , say s (regardless of its size k).

- If $s = 0$ then $\chi = \mathbf{0}$, clearly realizable in \mathbb{F}_2 .*
- Given χ with $|\text{supp}(\chi)| = s$ and being Howson, define the *cone* of χ with vertex $I \subseteq [k]$, denoted by $c_I(\chi)$, as*

$$c_I(\chi): \mathcal{P}([k]) \setminus \{\emptyset\} \rightarrow \{0, 1\}$$

$$J \mapsto \begin{cases} 0 & \text{if } J \not\subseteq I, \\ (J)\chi & \text{if } J \subseteq I. \end{cases}$$

- Now let $I_1, \dots, I_p \subseteq [k]$ be the maximal elements in $\text{supp}(\chi)$ (w.r.t. inclusion). It is clear that $\chi = c_{I_1}(\chi) \vee \dots \vee c_{I_p}(\chi)$.*

Characterization for the free case

Theorem (Delgado–Roy–V., '22)

A k -configuration is realizable in \mathbb{F}_n , $n \geq 2 \Leftrightarrow$ it is Howson.

(Proof)

For \Leftarrow , we will do induction on the cardinal of the support of χ , say s (regardless of its size k).

- If $s = 0$ then $\chi = \mathbf{0}$, clearly realizable in \mathbb{F}_2 .*
- Given χ with $|\text{supp}(\chi)| = s$ and being Howson, define the **cone of χ with vertex $I \subseteq [k]$** , denoted by $c_I(\chi)$, as*

$$c_I(\chi): \mathcal{P}([k]) \setminus \{\emptyset\} \rightarrow \{0, 1\}$$

$$J \mapsto \begin{cases} 0 & \text{if } J \not\subseteq I, \\ (J)\chi & \text{if } J \subseteq I. \end{cases}$$

- Now let $I_1, \dots, I_p \subseteq [k]$ be the maximal elements in $\text{supp}(\chi)$ (w.r.t. inclusion). It is clear that $\chi = c_{I_1}(\chi) \vee \dots \vee c_{I_p}(\chi)$.*

Characterization for the free case

Theorem (Delgado–Roy–V., '22)

A k -configuration is realizable in \mathbb{F}_n , $n \geq 2 \Leftrightarrow$ it is Howson.

(Proof)

For \Leftarrow , we will do induction on the cardinal of the support of χ , say s (regardless of its size k).

- If $s = 0$ then $\chi = \mathbf{0}$, clearly realizable in \mathbb{F}_2 .*
- Given χ with $|\text{supp}(\chi)| = s$ and being Howson, define the **cone of χ with vertex $I \subseteq [k]$** , denoted by $c_I(\chi)$, as*

$$c_I(\chi): \mathcal{P}([k]) \setminus \{\emptyset\} \rightarrow \{0, 1\}$$

$$J \mapsto \begin{cases} 0 & \text{if } J \not\subseteq I, \\ (J)\chi & \text{if } J \subseteq I. \end{cases}$$

- Now let $I_1, \dots, I_p \subseteq [k]$ be the maximal elements in $\text{supp}(\chi)$ (w.r.t. inclusion). It is clear that $\chi = c_{I_1}(\chi) \vee \dots \vee c_{I_p}(\chi)$.*

Characterization for the free case

(cont.)

- If $p \geq 2$, by the induction hypothesis we can realize each of $c_{I_1}(\chi), \dots, c_{I_p}(\chi)$ in \mathbb{F}_2 , and so, realize their join χ , in \mathbb{F}_2 as well.
- Hence, we are reduced to the case $p = 1$: χ is Howson and $\exists \emptyset \neq I_1 \subseteq [k]$ with $(I_1)\chi = 1$, and $(J)\chi = 0$ for every $J \not\subseteq I_1$.
- If $I_1 \neq [k]$ then any $j \in [k] \setminus I_1$ is 0-monochromatic, $\chi = \chi_{[j]} \boxplus_0 \mathbf{0}$, and we are reduced to realize $\chi_{[j]}$; repeating, we can assume $I_1 = [k]$. That is, χ is a Howson k -config. s.t. $([k])\chi = 1$.
- If $\chi = \mathbf{1}$ then it is clearly realizable in \mathbb{F}_2 .
- Otherwise, take $\emptyset \neq I_2 \subseteq [k]$ with $(I_2)\chi = 0$ and with maximal possible cardinal.
- Since $I_2 \neq [k]$, $\exists j \notin I_2$, and any such index is 1-monochromatic: in fact, any $J \subseteq [k]$ satisfies $|I_2 \cup J| > |I_2|$ so $(I_2 \cup J)\chi = 1$ and, since χ is Howson and $(I_2)\chi = 0$, then $(J)\chi = 1$.
- Hence, by induction hypothesis, $\chi_{[j]}$ is realizable in \mathbb{F}_2 and $\chi = \chi_{[j]} \boxplus_1 \mathbf{1}$ as well.



Characterization for the free case

(cont.)

- If $p \geq 2$, by the induction hypothesis we can realize each of $c_{I_1}(\chi), \dots, c_{I_p}(\chi)$ in \mathbb{F}_2 , and so, realize their join χ , in \mathbb{F}_2 as well.
- Hence, we are reduced to the case $p = 1$: χ is Howson and $\exists \emptyset \neq I_1 \subseteq [k]$ with $(I_1)\chi = 1$, and $(J)\chi = 0$ for every $J \not\subseteq I_1$.
- If $I_1 \neq [k]$ then any $j \in [k] \setminus I_1$ is 0-monochromatic, $\chi = \chi_{[j]} \boxplus_0 \mathbf{0}$, and we are reduced to realize $\chi_{[j]}$; repeating, we can assume $I_1 = [k]$. That is, χ is a Howson k -config. s.t. $([k])\chi = 1$.
- If $\chi = \mathbf{1}$ then it is clearly realizable in \mathbb{F}_2 .
- Otherwise, take $\emptyset \neq I_2 \subseteq [k]$ with $(I_2)\chi = 0$ and with maximal possible cardinal.
- Since $I_2 \neq [k]$, $\exists j \notin I_2$, and any such index is 1-monochromatic: in fact, any $J \subseteq [k]$ satisfies $|I_2 \cup J| > |I_2|$ so $(I_2 \cup J)\chi = 1$ and, since χ is Howson and $(I_2)\chi = 0$, then $(J)\chi = 1$.
- Hence, by induction hypothesis, $\chi_{[j]}$ is realizable in \mathbb{F}_2 and $\chi = \chi_{[j]} \boxplus_1 \mathbf{1}$ as well.



Characterization for the free case

(cont.)

- If $p \geq 2$, by the induction hypothesis we can realize each of $c_{I_1}(\chi), \dots, c_{I_p}(\chi)$ in \mathbb{F}_2 , and so, realize their join χ , in \mathbb{F}_2 as well.
- Hence, we are reduced to the case $p = 1$: χ is Howson and $\exists \emptyset \neq I_1 \subseteq [k]$ with $(I_1)\chi = 1$, and $(J)\chi = 0$ for every $J \not\subseteq I_1$.
- If $I_1 \neq [k]$ then any $j \in [k] \setminus I_1$ is 0-monochromatic, $\chi = \chi_{[j]} \boxplus_0 \mathbf{0}$, and we are reduced to realize $\chi_{[j]}$; repeating, we can assume $I_1 = [k]$. That is, χ is a Howson k -config. s.t. $([k])\chi = 1$.
- If $\chi = \mathbf{1}$ then it is clearly realizable in \mathbb{F}_2 .
- Otherwise, take $\emptyset \neq I_2 \subseteq [k]$ with $(I_2)\chi = 0$ and with maximal possible cardinal.
- Since $I_2 \neq [k]$, $\exists j \notin I_2$, and any such index is 1-monochromatic: in fact, any $J \subseteq [k]$ satisfies $|I_2 \cup J| > |I_2|$ so $(I_2 \cup J)\chi = 1$ and, since χ is Howson and $(I_2)\chi = 0$, then $(J)\chi = 1$.
- Hence, by induction hypothesis, $\chi_{[j]}$ is realizable in \mathbb{F}_2 and $\chi = \chi_{[j]} \boxplus_1 \mathbf{1}$ as well.



Characterization for the free case

(cont.)

- If $p \geq 2$, by the induction hypothesis we can realize each of $c_{I_1}(\chi), \dots, c_{I_p}(\chi)$ in \mathbb{F}_2 , and so, realize their join χ , in \mathbb{F}_2 as well.
- Hence, we are reduced to the case $p = 1$: χ is Howson and $\exists \emptyset \neq I_1 \subseteq [k]$ with $(I_1)\chi = 1$, and $(J)\chi = 0$ for every $J \not\subseteq I_1$.
- If $I_1 \neq [k]$ then any $j \in [k] \setminus I_1$ is 0-monochromatic, $\chi = \chi_{[j]} \boxplus_0 \mathbf{0}$, and we are reduced to realize $\chi_{[j]}$; repeating, we can assume $I_1 = [k]$. That is, χ is a Howson k -config. s.t. $([k])\chi = 1$.
- If $\chi = \mathbf{1}$ then it is clearly realizable in \mathbb{F}_2 .
- Otherwise, take $\emptyset \neq I_2 \subseteq [k]$ with $(I_2)\chi = 0$ and with maximal possible cardinal.
- Since $I_2 \neq [k]$, $\exists j \notin I_2$, and any such index is 1-monochromatic: in fact, any $J \subseteq [k]$ satisfies $|I_2 \cup J| > |I_2|$ so $(I_2 \cup J)\chi = 1$ and, since χ is Howson and $(I_2)\chi = 0$, then $(J)\chi = 1$.
- Hence, by induction hypothesis, $\chi_{[j]}$ is realizable in \mathbb{F}_2 and $\chi = \chi_{[j]} \boxplus_1 \mathbf{1}$ as well.



Characterization for the free case

(cont.)

- If $p \geq 2$, by the induction hypothesis we can realize each of $c_{I_1}(\chi), \dots, c_{I_p}(\chi)$ in \mathbb{F}_2 , and so, realize their join χ , in \mathbb{F}_2 as well.
- Hence, we are reduced to the case $p = 1$: χ is Howson and $\exists \emptyset \neq I_1 \subseteq [k]$ with $(I_1)\chi = 1$, and $(J)\chi = 0$ for every $J \not\subseteq I_1$.
- If $I_1 \neq [k]$ then any $j \in [k] \setminus I_1$ is 0-monochromatic, $\chi = \chi_{[j]} \boxplus_0 \mathbf{0}$, and we are reduced to realize $\chi_{[j]}$; repeating, we can assume $I_1 = [k]$. That is, χ is a Howson k -config. s.t. $([k])\chi = 1$.
- If $\chi = \mathbf{1}$ then it is clearly realizable in \mathbb{F}_2 .
- Otherwise, take $\emptyset \neq I_2 \subseteq [k]$ with $(I_2)\chi = 0$ and with maximal possible cardinal.
- Since $I_2 \neq [k]$, $\exists j \notin I_2$, and any such index is 1-monochromatic: in fact, any $J \subseteq [k]$ satisfies $|I_2 \cup J| > |I_2|$ so $(I_2 \cup J)\chi = 1$ and, since χ is Howson and $(I_2)\chi = 0$, then $(J)\chi = 1$.
- Hence, by induction hypothesis, $\chi_{[j]}$ is realizable in \mathbb{F}_2 and $\chi = \chi_{[j]} \boxplus_1 \mathbf{1}$ as well.



Characterization for the free case

(cont.)

- If $p \geq 2$, by the induction hypothesis we can realize each of $c_{I_1}(\chi), \dots, c_{I_p}(\chi)$ in \mathbb{F}_2 , and so, realize their join χ , in \mathbb{F}_2 as well.
- Hence, we are reduced to the case $p = 1$: χ is Howson and $\exists \emptyset \neq I_1 \subseteq [k]$ with $(I_1)\chi = 1$, and $(J)\chi = 0$ for every $J \not\subseteq I_1$.
- If $I_1 \neq [k]$ then any $j \in [k] \setminus I_1$ is 0-monochromatic, $\chi = \chi_{[j]} \boxplus_0 \mathbf{0}$, and we are reduced to realize $\chi_{[j]}$; repeating, we can assume $I_1 = [k]$. That is, χ is a Howson k -config. s.t. $([k])\chi = 1$.
- If $\chi = \mathbf{1}$ then it is clearly realizable in \mathbb{F}_2 .
- Otherwise, take $\emptyset \neq I_2 \subseteq [k]$ with $(I_2)\chi = 0$ and with maximal possible cardinal.
- Since $I_2 \neq [k]$, $\exists j \notin I_2$, and any such index is 1-monochromatic: in fact, any $J \subseteq [k]$ satisfies $|I_2 \cup J| > |I_2|$ so $(I_2 \cup J)\chi = 1$ and, since χ is Howson and $(I_2)\chi = 0$, then $(J)\chi = 1$.
- Hence, by induction hypothesis, $\chi_{[j]}$ is realizable in \mathbb{F}_2 and $\chi = \chi_{[j]} \boxplus_1 \mathbf{1}$ as well.



Characterization for the free case

(cont.)

- If $p \geq 2$, by the induction hypothesis we can realize each of $c_{I_1}(\chi), \dots, c_{I_p}(\chi)$ in \mathbb{F}_2 , and so, realize their join χ , in \mathbb{F}_2 as well.
- Hence, we are reduced to the case $p = 1$: χ is Howson and $\exists \emptyset \neq I_1 \subseteq [k]$ with $(I_1)\chi = 1$, and $(J)\chi = 0$ for every $J \not\subseteq I_1$.
- If $I_1 \neq [k]$ then any $j \in [k] \setminus I_1$ is 0-monochromatic, $\chi = \chi_{[j]} \boxplus_0 \mathbf{0}$, and we are reduced to realize $\chi_{[j]}$; repeating, we can assume $I_1 = [k]$. That is, χ is a Howson k -config. s.t. $([k])\chi = 1$.
- If $\chi = \mathbf{1}$ then it is clearly realizable in \mathbb{F}_2 .
- Otherwise, take $\emptyset \neq I_2 \subseteq [k]$ with $(I_2)\chi = 0$ and with maximal possible cardinal.
- Since $I_2 \neq [k]$, $\exists j \notin I_2$, and any such index is 1-monochromatic: in fact, any $J \subseteq [k]$ satisfies $|I_2 \cup J| > |I_2|$ so $(I_2 \cup J)\chi = 1$ and, since χ is Howson and $(I_2)\chi = 0$, then $(J)\chi = 1$.
- Hence, by induction hypothesis, $\chi_{[j]}$ is realizable in \mathbb{F}_2 and $\chi = \chi_{[j]} \boxplus_1 \mathbf{1}$ as well.



Outline

- 1 Our main results
- 2 Free-times-free-abelian groups
- 3 Realizable / unrealizable k -configurations
- 4 The free case
- 5 Open questions

Open questions

Question

Can we characterize the k -configurations realizable in $\mathbb{F}_n \times \mathbb{Z}^m$, for each particular m ?

Question

Is there an algorithm which, on input m and χ , decides whether χ is realizable in $\mathbb{F}_n \times \mathbb{Z}^m$ (and, in the affirmative case, computes such a realization)?

Question

Is there a finitely presented intersection-saturated group G which does not contain $\mathbb{F}_2 \times \mathbb{Z}^m$, for some $m \in \mathbb{N}$?

Open questions

Question

Can we characterize the k -configurations realizable in $\mathbb{F}_n \times \mathbb{Z}^m$, for each particular m ?

Question

Is there an algorithm which, on input m and χ , decides whether χ is realizable in $\mathbb{F}_n \times \mathbb{Z}^m$ (and, in the affirmative case, computes such a realization)?

Question

Is there a finitely presented intersection-saturated group G which does not contain $\mathbb{F}_2 \times \mathbb{Z}^m$, for some $m \in \mathbb{N}$?

Open questions

Question

Can we characterize the k -configurations realizable in $\mathbb{F}_n \times \mathbb{Z}^m$, for each particular m ?

Question

Is there an algorithm which, on input m and χ , decides whether χ is realizable in $\mathbb{F}_n \times \mathbb{Z}^m$ (and, in the affirmative case, computes such a realization)?

Question

Is there a finitely presented intersection-saturated group G which does not contain $\mathbb{F}_2 \times \mathbb{Z}^m$, for some $m \in \mathbb{N}$?

DZIĘKUJĘ

THANKS